


Exercise - 12A

1.

Sol:



Let O be the center of the given circle.

Let P be a point, such that

$$OP = 17 \text{ cm.}$$

Let OT be the radius, where

$$OT = 5 \text{ cm}$$

Join TP , where TP is a tangent.

Now, tangent drawn from an external point is perpendicular to the radius at the point of contact.

$$\therefore OT \perp TP$$

In the right $\triangle OTP$, we have:

$$OP^2 = OT^2 + TP^2 \quad [\text{By Pythagoras' theorem:}]$$

$$TP = \sqrt{OP^2 - OT^2}$$

$$= \sqrt{17^2 - 5^2}$$

$$= \sqrt{289 - 25}$$


$$= \sqrt{264}$$

$$= 16.25 \text{ cm}$$

\therefore The length of the tangent is 16.25 cm.

2.

Sol:



Draw a circle and let P be a point such that $OP = 25 \text{ cm}$.

Let TP be the tangent, so that $TP = 24 \text{ cm}$

Join OT where OT is radius.

Now, tangent drawn from an external point is perpendicular to the radius at the point of contact.

$$\therefore OT \perp PT$$

In the right $\triangle OTP$, we have:

$$OP^2 = OT^2 + TP^2 \quad [\text{By Pythagoras' theorem:}]$$

$$OT^2 = \sqrt{OP^2 - TP^2}$$

$$= \sqrt{25^2 - 24^2}$$

$$= \sqrt{625 - 576}$$


$$= \sqrt{49}$$

$$= 7 \text{ cm}$$

\therefore The length of the radius is 7cm.

3.

Sol:



We know that the radius and tangent are perpendicular at their point of contact

In right triangle AOP

$$AO^2 = OP^2 + PA^2$$

$$\Rightarrow (6.5)^2 = (2.5)^2 + PA^2$$

$$\Rightarrow PA^2 = 36$$

$$\Rightarrow PA = 6 \text{ cm}$$


Since, the perpendicular drawn from the center bisects the chord.

$$\therefore PA = PB = 6 \text{ cm}$$

Now, $AB = AP + PB = 6 + 6 = 12 \text{ cm}$

Hence, the length of the chord of the larger circle is 12cm.

4.



Sol:

We know that tangent segments to a circle from the same external point are congruent.

Now, we have

$$AD = AF, BD = BE \text{ and } CE = CF$$

$$\text{Now, } AD + BD = 12\text{cm} \quad \dots\dots(1)$$

$$AF + FC = 10 \text{ cm} \\ \Rightarrow AD + FC = 10 \text{ cm} \quad \dots\dots(2)$$

$$BE + EC = 8 \text{ cm} \\ \Rightarrow BD + FC = 8\text{cm} \quad \dots\dots(3)$$

Adding all these we get

$$AD + BD + AD + FC + BD + FC = 30 \\ \Rightarrow 2(AD + BD + FC) = 30 \\ \Rightarrow AD + BD + FC = 15\text{cm} \quad \dots\dots(4)$$

Solving (1) and (4), we get

$$FC = 3 \text{ cm}$$

Solving (2) and (4), we get

$$BD = 5 \text{ cm}$$


Solving (3) and (4), we get

$$\text{and } AD = 7 \text{ cm}$$

$$\therefore AD = AF = 7 \text{ cm, } BD = BE = 5 \text{ cm and } CE = CF = 3 \text{ cm}$$

5.

Sol:



Let the circle touch the sides of the quadrilateral AB , BC , CD and DA at P , Q , R and S respectively.

Given, $AB = 6\text{cm}$, $BC = 7\text{ cm}$ and $CD = 4\text{cm}$.

Tangents drawn from an external point are equal.

$$AP = AS, BP = BQ, CR = CQ \text{ and } DR = DS$$

Now, $AB + CD = (AP + BP) + (CR + DR)$

$$\Rightarrow AB + CD = (AS + BQ) + (CQ + DS)$$

$$\Rightarrow AB + CD = (AS + DS) + (BQ + CQ)$$

$$\Rightarrow AB + CD = AD + BC$$

$$\Rightarrow AD = (AB + CD) - BC$$

$$\Rightarrow AD = (6 + 4) - 7$$


$$\Rightarrow AD = 3\text{ cm.}$$

\therefore The length of AD is 3 cm.

6.

Sol:

Construction: Join OA , OC and OB



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OCA = \angle OCB = 90^\circ$$

Now, In $\triangle OCA$ and $\triangle OCB$

$$\angle OCA = \angle OCB = 90^\circ$$

$$OA = OB \text{ (Radii of the larger circle)}$$


$$OC = OC \text{ (Common)}$$

By RHS congruency

$$\triangle OCA \cong \triangle OCB$$

$$\therefore CA = CB$$

7.

**Sol:**

Given, PA and PB are the tangents to a circle with center O and CD is a tangent at E and PA = 14 cm.

Tangents drawn from an external point are equal.

$$\therefore PA = PB, CA = CE \text{ and } DB = DE$$

$$\text{Perimeter of } \triangle PCD = PC + CD + PD$$

$$= (PA - CA) + (CE + DE) + (PB - DB)$$

$$= (PA - CE) + (CE + DE) + (PB - DE)$$

$$= (PA + PB)$$

$$= 2PA \quad (\because PA = PB)$$


$$= (2 \times 14) \text{ cm}$$

$$= 28 \text{ cm}$$

$$= 28 \text{ cm}$$

$$\therefore \text{Perimeter of } \triangle PCD = 28 \text{ cm.}$$

8.

**Sol:**

Given, a circle inscribed in triangle ABC, such that the circle touches the sides of the triangle

Tangents drawn to a circle from an external point are equal.

$$\therefore AP = AR = 7 \text{ cm}, CQ = CR = 5 \text{ cm.}$$

$$\text{Now, } BP = (AB - AP) = (10 - 7) = 3 \text{ cm}$$

$$\therefore BP = BQ = 3 \text{ cm}$$


$$\therefore BC = (BQ + QC)$$

$$\Rightarrow BC = 3 + 5$$

$$\Rightarrow BC = 8$$

\therefore The length of BC is 8 cm.

9.



Sol:

Here, $OA = OB$

And $OA \perp AP$, $OB \perp BP$ (Since tangents drawn from an external point are perpendicular to the radius at the point of contact)

$$\therefore \angle OAP = 90^\circ, \angle OBP = 90^\circ$$


$$\therefore \angle OAP + \angle OBP = 90^\circ + 90^\circ = 180^\circ$$

$$\therefore \angle AOB + \angle APB = 180^\circ \text{ (Since, } \angle OAP + \angle OBP + \angle AOB + \angle APB = 360^\circ \text{)}$$

Sum of opposite angle of a quadrilateral is 180° .

Hence A, O, B and P are concyclic.

10.



Sol:

We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$AR = AO, BR = BP \text{ and } CP = CQ$$

Now, $AB = AC$

$$\Rightarrow AR + RB = AQ + QC$$


$$\Rightarrow AR + RB = AR + OC$$

$$\Rightarrow RB = QC$$

$$\Rightarrow BP = CP$$

Hence, P bisects BC at P .

11.



Sol:

Given, O is the center of two concentric circles of radii $OA = 6$ cm and $OB = 4$ cm.
 PA and PB are the two tangents to the outer and inner circles respectively and $PA = 10$ cm.

Now, tangent drawn from an external point is perpendicular to the radius at the point of contact.

$$\therefore \angle OAP = \angle OBP = 90^\circ$$

$$\therefore \text{From right-angled } \triangle OAP, OP^2 = OA^2 + PA^2$$

$$\Rightarrow OP = \sqrt{OA^2 + PA^2}$$

$$\Rightarrow OP = \sqrt{6^2 + 10^2}$$

$$\Rightarrow OP = \sqrt{136} \text{ cm.}$$

$$\therefore \text{From right-angled } \triangle OBP, OP^2 = OB^2 + PB^2$$

$$\Rightarrow PB = \sqrt{OP^2 - OB^2}$$


$$\Rightarrow PB = \sqrt{136 - 16}$$

$$\Rightarrow PB = \sqrt{120} \text{ cm}$$

$$\Rightarrow PB = 10.9 \text{ cm.}$$


\therefore The length of PB is 10.9 cm.

12.



Sol:

Construction: Join $OA, OB, OC, OE \perp AB$ at E and $OF \perp AC$ at F



We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$AE = AF, BD = BE = 6 \text{ cm and } CD = CF = 9 \text{ cm}$$

Now,

$$\text{Area}(\triangle ABC) = \text{Area}(\triangle BOC) + \text{Area}(\triangle AOB) + \text{Area}(\triangle AOC)$$

$$\Rightarrow 54 = \frac{1}{2} \times BC \times OD + \frac{1}{2} \times AB \times OE + \frac{1}{2} \times AC \times OF$$

$$\Rightarrow 108 = 15 \times 3 + (6+x) \times 3 + (9+x) \times 3$$

$$\Rightarrow 36 = 15 + 6 + x + 9 + x$$

$$\Rightarrow 36 = 30 + 2x$$


$$\Rightarrow 2x = 6$$

$$\Rightarrow x = 3 \text{ cm}$$

$$\therefore AB = 6 + 3 = 9 \text{ cm and } AC = 9 + 3 = 12 \text{ cm}$$

13.

Sol:



Let $TR = y$ and $TP = x$

We know that the perpendicular drawn from the center to the chord bisects it.

$$\therefore PR = RQ$$

Now, $PR + RQ = 4.8$

$$\Rightarrow PR + PR = 4.8$$

$$\Rightarrow PR = 2.4$$

Now, in right triangle POR

By Using Pythagoras theorem, we have

$$PO^2 = OR^2 + PR^2$$

$$\Rightarrow 3^2 = OR^2 + (2.4)^2$$

$$\Rightarrow OR^2 = 3.24$$

$$\Rightarrow OR = 1.8$$

Now, in right triangle TPR

By Using Pythagoras theorem, we have

$$TP^2 = TR^2 + PR^2$$

$$\Rightarrow x^2 = y^2 + (2.4)^2$$

$$\Rightarrow x^2 = y^2 + 5.76 \quad \dots\dots(1)$$

Again, In right triangle TPQ

By Using Pythagoras theorem, we have

$$TO^2 = TP^2 + PO^2$$

$$\Rightarrow (y+1.8)^2 = x^2 + 3^2$$

$$\Rightarrow y^2 + 3.6y + 3.24 = x^2 + 9$$

$$\Rightarrow y^2 + 3.6y = x^2 + 5.76 \quad \dots\dots(2)$$


Solving (1) and (2), we get

$$x = 4 \text{ cm and } y = 3.2 \text{ cm}$$

$$\therefore TP = 4 \text{ cm}$$

14.

Sol:



Suppose CD and AB are two parallel tangents of a circle with center O

Construction: Draw a line parallel to CD passing through O i.e. OP

We know that the radius and tangent are perpendicular at their point of contact.

$$\angle OQC = \angle ORA = 90^\circ$$

$$\text{Now, } \angle OQC + \angle POQ = 180^\circ \quad (\text{co-interior angles})$$

$$\Rightarrow \angle POQ = 180^\circ - 90^\circ = 90^\circ$$

$$\text{Similarly, Now, } \angle ORA + \angle POR = 180^\circ (\text{co-interior angles})$$


$$\Rightarrow \angle POQ = 180^\circ - 90^\circ = 90^\circ$$

$$\text{Now, } \angle POR + \angle POQ = 90^\circ + 90^\circ = 180^\circ$$

Since, $\angle POR$ and $\angle POQ$ are linear pair angles whose sum is 180°

Hence, QR is a straight line passing through center O.

15.



Sol:

We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$DS = DR, AR = AQ$$

$$\text{Now } AD = 23 \text{ cm}$$

$$\Rightarrow AR + RD = 23$$

$$\Rightarrow AR = 23 - RD$$

$$\Rightarrow AR = 23 - 5 \quad [\because DS = DR = 5]$$

$$\Rightarrow AR = 18 \text{ cm}$$

$$\text{Again, } AB = 29 \text{ cm}$$

$$\Rightarrow AQ + QB = 29$$

$$\Rightarrow QB = 29 - AQ$$

$$\Rightarrow QB = 29 - 18 \quad [\because AR = AQ = 18]$$

$$\Rightarrow QB = 11 \text{ cm}$$


Since all the angles in a quadrilateral BQOP are right angles and $OP = BQ$

Hence, BQOP is a square.

We know that all the sides of square are equal.

Therefore, $BQ = PO = 11 \text{ cm}$

16.

**Sol:**

AB is the chord passing through the center

So, AB is the diameter

Since, angle in a semicircle is a right angle

$$\therefore \angle APB = 90^\circ$$

By using alternate segment theorem

$$\text{We have } \angle APB = \angle PAT = 30^\circ$$

Now, in $\triangle APB$

$$\angle BAP + \angle APB + \angle ABP = 180^\circ \text{ (Angle sum property of triangle)}$$

$$\Rightarrow \angle BAP = 180^\circ - 90^\circ - 30^\circ = 60^\circ$$

Now, $\angle BAP = \angle APT + \angle PTA$ (Exterior angle property)

$$\Rightarrow 60^\circ = 30^\circ + \angle PTA$$

$$\Rightarrow \angle PTA = 60^\circ - 30^\circ = 30^\circ$$

We know that sides opposite to equal angles are equal

$$\therefore AP = AT$$

In right triangle ABP

$$\sin \angle ABP = \frac{AP}{BA}$$


$$\Rightarrow \sin 30^\circ = \frac{AT}{BA}$$

$$\Rightarrow \frac{1}{2} = \frac{AT}{BA}$$

$$\therefore BA : AT = 2 : 1$$

Exercise – 12B

1.



Sol:

We know that when a quadrilateral circumscribes a circle then sum of opposite sides is equal to the sum of other opposite sides.

$$\therefore AB + CD = AD + BC$$


$$\Rightarrow 6 + 8 = AD = 9$$

$$\Rightarrow AD = 5 \text{ cm}$$

2.

Sol:

Construction: Join OB



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, In quadrilateral AOBP

$$\angle AOB + \angle OBP + \angle APB + \angle OAP = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle AOB + 90^\circ + 50^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow 230^\circ + \angle BOC = 360^\circ$$

$$\Rightarrow \angle AOB = 130^\circ$$

Now, In isosceles triangle AOB

$$\angle AOB + \angle OAB + \angle OBA = 180^\circ \quad [\text{Angle sum property of a triangle}]$$


$$\Rightarrow 130^\circ + 2\angle OAB = 180^\circ \quad [\because \angle OAB = \angle OBA]$$

$$\Rightarrow \angle OAB = 25^\circ$$

3.

Sol:

Construction: Join OQ and OT



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OTP = \angle OQP = 90^\circ$$

Now, In quadrilateral OQPT

$$\angle QOT + \angle OTP + \angle OQP + \angle TPO = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle QOT + 90^\circ + 90^\circ + 70^\circ = 360^\circ$$


$$\Rightarrow 250^\circ + \angle QOT = 360^\circ$$

$$\Rightarrow \angle QOT = 110^\circ$$

We know that the angle subtended by an arc at the center is double the angle subtended by the arc at any point on the remaining part of the circle.

$$\therefore \angle TRQ = \frac{1}{2}(\angle QOT) = 55^\circ$$

4.



Sol:

We know that tangent segments to a circle from the same external point are congruent.

So, we have

$$EA = EC \text{ for the circle having center } O_1$$

and

$$ED = EB \text{ for the circle having center } O_2$$


Now, Adding ED on both sides in $EA = EC$. we get

$$EA + ED = EC + ED$$

$$\Rightarrow EA + EB = EC + ED$$

$$\Rightarrow AB = CD$$

5.



Sol:

We know that the radius and tangent are perpendicular at their point of contact.

$$\therefore \angle OPT = 90^\circ$$

$$\text{Now, } \angle OPQ = \angle OPT - \angle TPQ = 90^\circ - 70^\circ = 20^\circ$$

Since, $OP = OQ$ as both are radius

$$\therefore \angle OPQ = \angle OQP = 20^\circ \quad (\text{Angles opposite to equal sides are equal})$$

Now, In isosceles ΔPOQ


$$\angle POQ + \angle OPQ + \angle OQP = 180^\circ \quad (\text{Angle sum property of a triangle})$$

$$\Rightarrow \angle POQ = 180^\circ - 20^\circ = 140^\circ$$

6.

Sol:

Construction: Join $OA, OB, OC, OE \perp AB$ at E and $OF \perp AC$ at F



We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$AE = AF, BD = BE = 4\text{ cm and } CD = CF = 3\text{ cm}$$


Now,

$$\text{Area}(\Delta ABC) = \text{Area}(\Delta BOC) + \text{Area}(\Delta AOB) + \text{Area}(\Delta AOC)$$

$$\begin{aligned} \Rightarrow 21 &= \frac{1}{2} \times BC \times OD + \frac{1}{2} \times AB \times OE + \frac{1}{2} \times AC \times OF \\ \Rightarrow 42 &= 7 \times 2 + (4+x) \times 2 + (3+x) \times 2 \\ \Rightarrow 21 &= 7 + 4 + x + 3 + x \\ \Rightarrow 21 &= 14 + 2x \\ \Rightarrow 2x &= 7 \\ \Rightarrow x &= 3.5 \text{ cm} \\ \therefore AB &= 4 + 3.5 = 7.5 \text{ cm and } AC = 3 + 3.5 = 6.5 \text{ cm} \end{aligned}$$

7.

Sol:



Given Two circles have the same center O and AB is a chord of the larger circle touching the smaller circle at C; also. $OA = 5 \text{ cm}$ and $OC = 3 \text{ cm}$

In $\triangle OAC$, $OA^2 = OC^2 + AC^2$

$$\therefore AC^2 = OA^2 - OC^2$$

$$\Rightarrow AC^2 = 5^2 - 3^2$$

$$\Rightarrow AC^2 = 25 - 9$$

$$\Rightarrow AC^2 = 16$$

$$\Rightarrow AC = 4 \text{ cm}$$


$\therefore AB = 2AC$ (Since perpendicular drawn from the center of the circle bisects the chord)

$$\therefore AB = 2 \times 4 = 8 \text{ cm}$$

The length of the chord of the larger circle is 8 cm.

8.

Sol:



Let AB be the tangent to the circle at point P with center O.

To prove: PQ passes through the point O.

Construction: Join OP.

Through O, draw a straight line CD parallel to the tangent AB.

Proof: Suppose that PQ doesn't pass through point O.

PQ intersect CD at R and also intersect AB at P

AS, $CD \parallel AB$. PQ is the line of intersection.

$\angle ORP = \angle RPA$ (Alternate interior angles)

but also.

$\angle RPA = 90^\circ$ ($OP \perp AB$)

$\Rightarrow \angle ORP = 90^\circ$

$\angle ROP + \angle OPA = 180^\circ$ (Co interior angles)

$\Rightarrow \angle ROP + 90^\circ = 180^\circ$

$\Rightarrow \angle ROP = 90^\circ$


Thus, the $\triangle ORP$ has 2 right angles i.e., $\angle ORP$ and $\angle ROP$ which is not possible

Hence, our supposition is wrong

\therefore PQ passes through the point O.

9.

Sol:



Construction Join PO and OQ

In $\triangle POR$ and $\triangle QOR$

$OP = OQ$ (Radii)

$RP = RQ$ (Tangents from the external point are congruent)

$OR = OR$ (Common)

By SSS congruency, $\triangle POR \cong \triangle QOR$

$\angle PRO = \angle QRO$ (C.P.C.T)

Now, $\angle PRO + \angle QRO = \angle PRQ$

$$\Rightarrow 2\angle PRO = 120^\circ$$

$$\Rightarrow \angle PRO = 60^\circ$$

Now. In $\triangle POR$

$$\cos 60^\circ = \frac{PR}{OR}$$


$$\Rightarrow \frac{1}{2} = \frac{PR}{OR}$$

$$\Rightarrow OR = 2PR$$

$$\Rightarrow OR = PR + PR$$

$$\Rightarrow OR = PR + RQ$$

10.



Sol:

We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$AD = AF, BD = BE \text{ and } CE = CF$$

$$\text{Now } AD + BD = 14 \text{ cm} \quad \dots\dots(1)$$

$$AF + FC = 12 \text{ cm}$$

$$\Rightarrow AD + FC = 12 \text{ cm} \quad \dots\dots(2)$$

$$BE + EC = 8 \text{ cm}$$

$$\Rightarrow BD + FC = 8 \text{ cm} \quad \dots\dots(3)$$

Adding all these we get

$$AD + BD + AD + FC + BD + FC = 34$$

$$\Rightarrow 2(AD + BD + FC) = 34$$

$$\Rightarrow AD + BO + FC = 17 \text{ cm} \quad \dots\dots\dots(4)$$

Solving (1) and (4), we get


$$FC = 3 \text{ cm}$$

Solving (2) and (4), we get

$$BD = 5 \text{ cm} = BE$$

Solving (3) and (4), we get
and $AD = 9 \text{ cm}$

11.



Sol:

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, In quadrilateral AOBP

$$\angle APB + \angle AOB + \angle OBP + \angle OAP = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle APB + \angle AOB + 90^\circ + 90^\circ = 360^\circ$$


$$\Rightarrow \angle APB + \angle AOB = 180^\circ$$

Since, the sum of the opposite angles of the quadrilateral is 180°

Hence, AOBP is a cyclic quadrilateral

12.

Sol:



We know that the radius and tangent are perpendicular at their point of contact

Since, the perpendicular drawn from the centre bisect the chord

$$\therefore AP = PB = \frac{AB}{2} = 4 \text{ cm}$$

In right triangle AOP

$$AO^2 = OP^2 + PA^2$$


$$\Rightarrow 5^2 = OP^2 + 4^2$$

$$\Rightarrow OP^2 = 9$$

$$\Rightarrow OP = 3 \text{ cm}$$

Hence, the radius of the smaller circle is 3 cm.

13.



Sol:

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OPT = 90^\circ$$

$$\text{Now, } \angle OPQ = \angle OPT - \angle QPT = 90^\circ - 60^\circ = 30^\circ$$

Since, $OP = OQ$ as both are radii

$$\therefore \angle OPQ = \angle OQP = 30^\circ \text{ (Angles opposite to equal sides are equal)}$$

Now, In isosceles, POQ

$$\angle POQ + \angle OPQ + \angle OQP = 180^\circ \text{ (Angle sum property of a triangle)}$$

$$\Rightarrow \angle POQ = 180^\circ - 30^\circ - 30^\circ = 120^\circ$$


Now, $\angle POQ + \text{reflex } \angle POQ = 360^\circ$ (Complete angle)

$$\Rightarrow \text{reflex } \angle POQ = 360^\circ - 120^\circ = 240^\circ$$

We know that the angle subtended by an arc at the centre double the angle subtended by the arc at any point on the remaining part of the circle


$$\therefore \angle PRQ = \frac{1}{2}(\text{reflex } \angle POQ) = 120^\circ$$

14.



Sol:

Construction: Join OB



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, In quadrilateral AOBP

$$\angle AOB + \angle OBP + \angle APB + \angle OAP = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle AOB + 90^\circ + 60^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow 240^\circ + \angle AOB = 360^\circ$$

$$\Rightarrow \angle AOB = 120^\circ$$

Now, In isosceles triangle AOB

$$\angle AOB + \angle OAB + \angle OBA = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 120^\circ + 2\angle OAB = 180^\circ \quad [\because \angle OAB = \angle OBA]$$

$$\Rightarrow \angle OAB = 30^\circ$$


Exercise – Multiple Choice Questions

1.


Answer: (b) 2

Sol:

We can draw only two tangents from an external point to a circle.



2.



Answer: (c) 5 cm

Sol:

We know that the radius and tangent are perpendicular at their point of contact


$$OQ = \frac{1}{2}QS = 3\text{ cm} \quad [\because \text{Radius is half of diameter}]$$

Now, in right triangle OQR

By using Pythagoras theorem, we have

$$\begin{aligned}
 OR^2 &= RQ^2 + OQ^2 \\
 &= 4^2 + 3^2 \\
 &= 16 + 9 \\
 &= 25 \\
 \therefore OR^2 &= 25 \\
 \Rightarrow OR &= 5 \text{ cm}
 \end{aligned}$$

3.

**Answer:** (c) 25 cm**Sol:**

The tangent at any point of a circle is perpendicular to the radius at the point of contact

$$\therefore OT \perp PT$$

From right – angled triangle PTO ,

$$\therefore OP^2 = OT^2 + PT^2 \quad [\text{Using Pythagoras' theorem}]$$

$$\Rightarrow OP^2 = (7)^2 + (24)^2$$

$$\Rightarrow OP^2 = 49 + 576$$

$$\Rightarrow OP^2 = 625$$

$$\Rightarrow OP = \sqrt{625}$$

$$\Rightarrow OP = 25 \text{ cm}$$


4.

Answer: (d) two diameters**Sol:**

Two diameters cannot be parallel as they perpendicularly bisect each other.

5.

Answer: (c) $10\sqrt{2}$

Sol:

In right triangle AOB

By using Pythagoras theorem, we have

$$AB^2 = BO^2 + OA^2$$


$$= 10^2 + 10^2$$

$$= 100 + 100$$

$$= 200$$

$$\therefore OR^2 = 200$$

$$\Rightarrow OR = 10\sqrt{2} \text{ cm}$$

6.**Answer:** (a) 8 cm**Sol:**

In right triangle PTO

By using Pythagoras theorem, we have

$$PO^2 = OT^2 + TP^2$$

$$\Rightarrow 10^2 = 6^2 + TP^2$$

$$\Rightarrow 100 = 36 + TP^2$$


$$\Rightarrow TP^2 = 64$$

$$\Rightarrow TP = 8 \text{ cm}$$

7.

Answer: (a) 10 cm**Sol:**

Construction: Join OT.



We know that the radius and tangent are perpendicular at their point of contact

In right triangle PTO

By using Pythagoras theorem, we have

$$PO^2 = OT^2 + TP^2$$


$$\Rightarrow 26^2 = OT^2 + 24^2$$

$$\Rightarrow 676 = OT^2 + 576$$

$$\Rightarrow TP^2 = 100$$

$$\Rightarrow TP = 10 \text{ cm}$$

8.

Answer: 45° **Sol:**

We know that the radius and tangent are perpendicular at their point of contact


Now, In isosceles right triangle POQ

$$\angle POQ + \angle OPQ + \angle OQP = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 2\angle OQP + 90^\circ = 180^\circ$$

$$\Rightarrow \angle OQP = 45^\circ$$

9.

**Answer: (d) 140°** **Sol:**

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBA = \angle OCA = 90^\circ$$

Now, In quadrilateral ABOC


$$\angle BAC + \angle OCA + \angle OBA + \angle BOC = 360^\circ \quad [\text{Angle sum property of quadrilateral}]$$

$$\Rightarrow 40^\circ + 90^\circ + 90^\circ + \angle BOC = 360^\circ$$

$$\Rightarrow 220^\circ + \angle BOC = 360^\circ$$

$$\Rightarrow \angle BOC = 140^\circ$$

10.

Answer: (d) 120° **Sol:**

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBC = \angle OAC = 90^\circ$$

Now, In quadrilateral ABOC


$$\angle ACB + \angle OAC + \angle OBC + \angle AOB = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle ACB + 90^\circ + 90^\circ + 60^\circ = 360^\circ$$

$$\Rightarrow \angle ACB + 240^\circ = 360^\circ$$

$$\Rightarrow \angle ACB = 120^\circ$$

11.

**Answer:** (c) 16 cm**Sol:**

We know that the radius and tangent are perpendicular at their point of contact

In right triangle AOP

$$AO^2 = OP^2 + PA^2$$

$$\Rightarrow 10^2 = 6^2 + PA^2$$

$$\Rightarrow PA^2 = 64$$


$$\Rightarrow PA = 8 \text{ cm}$$

Since, the perpendicular drawn from the center bisect the chord

$$\therefore PA = PB = 8 \text{ cm}$$

Now, $AB = AP + PB = 8 + 8 = 16 \text{ cm}$

12.

**Answer:** (b) 15**Sol:**

We know that the radius and tangent are perpendicular at their point of contact

In right triangle AOB

By using Pythagoras theorem, we have

$$OA^2 = AB^2 + OB^2$$

$$\Rightarrow 17^2 = AB^2 + 8^2$$

$$\Rightarrow 289 = AB^2 + 64$$


$$\Rightarrow AB^2 = 225$$

$$\Rightarrow AB = 15 \text{ cm}$$

The tangents drawn from the external point are equal

Therefore, the length of AC is 15 cm

13.



Answer: (b) 50°

Sol:

$\angle ABC = 90^\circ$ (Angle in a semicircle)

In $\triangle ABC$, we have: $\angle ACB + \angle CAB + \angle ABC = 180^\circ$

$$\Rightarrow 50^\circ + \angle CAB + 90^\circ = 180^\circ$$

$$\Rightarrow \angle CAB = (180^\circ - 140^\circ)$$

$$\Rightarrow \angle CAB = 40^\circ$$

Now, $\angle CAT = 90^\circ$ (Tangents drawn from an external point are perpendicular to the radius at the point of contact)


$$\therefore \angle CAB + \angle BAT = 90^\circ$$

$$\Rightarrow 40^\circ + \angle BAT = 90^\circ$$

$$\Rightarrow \angle BAT = (90^\circ - 40^\circ)$$

$$\Rightarrow \angle BAT = 50^\circ$$

14.



Answer: (a) 35°

Sol:

We know that the radius and tangent are perpendicular at their point of contact

Since, $OP = OQ$

$\therefore POQ$ is a isosceles right triangle

Now, In isosceles right triangle POQ

$$\angle POQ + \angle OPQ + \angle OQP = 180^\circ \quad [\text{Angle sum proper of a triangle}]$$


$$\Rightarrow 70^\circ + 2\angle OPQ = 180^\circ$$

$$\Rightarrow \angle OPQ = 55^\circ$$

Now, $\angle TPQ + \angle OPQ = 90^\circ$

$$\Rightarrow \angle TPQ = 35^\circ$$

15.



Answer: (c) $2\sqrt{3}$ cm

Sol:

$$OA \perp AT$$


$$\text{So, } \frac{AT}{OT} = \cos 30^\circ$$

$$\Rightarrow \frac{AT}{4} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow AT = \left(\frac{\sqrt{3}}{2} \times 4 \right)$$

$$\Rightarrow AT = 2\sqrt{3}$$

16.



Answer: (c) 70°

Sol:

Given, PA and PB are tangents to a circle with center O, with $\angle AOB = 110^\circ$.

Now, we know that tangents drawn from an external point are perpendicular to the radius at the point of contact.

So, $\angle OAP = 90^\circ$ and $\angle OBP = 90^\circ$

$\Rightarrow \angle OAP + \angle OBP = 90^\circ + 90^\circ = 180^\circ$, which shows that OABP is a cyclic quadrilateral.


$$\therefore \angle AOB + \angle APB = 180^\circ$$

$$\Rightarrow 110^\circ + \angle APB = 180^\circ$$

$$\Rightarrow \angle APB = 180^\circ - 110^\circ$$

$$\Rightarrow \angle APB = 70^\circ$$

17.

**Answer:** (b) 10 cm**Sol:**

We know that tangent segments to a circle from the same external point are congruent

Therefore, we have

$$AF = AE = 4 \text{ cm}$$


$$BF = BD = 3 \text{ cm}$$

$$EC = AC - AE = 11 - 4 = 7 \text{ cm}$$

$$CD = CE = 7 \text{ cm}$$

$$\therefore BC = BD + DC = 3 + 7 = 10 \text{ cm}$$

18.


**Answer:** (b) 45° **Sol:**

We know that the sum of angles subtended by opposite sides of a quadrilateral having a circumscribed circle is 180 degrees

$$\therefore \angle AOD + \angle BOC = 180^\circ$$

$$\Rightarrow \angle BOC = 180^\circ - 135^\circ = 45^\circ$$

19.



Answer: (a) 100°

Sol:

Given, $\angle QPT = 50^\circ$

And $\angle OPT = 90^\circ$ (Tangents drawn from an external point are perpendicular to the radius at the point of contact)

$$\therefore \angle OPQ = (\angle OPT - \angle QPT) = (90^\circ - 50^\circ) = 40^\circ$$


$OP = OQ$ (Radius of the same circle)

$$\Rightarrow \angle OQP = \angle OPQ = 40^\circ$$

In $\triangle POQ$, $\angle POQ + \angle OQP + \angle OPQ = 180^\circ$

$$\therefore \angle POQ = 180^\circ - (40^\circ + 40^\circ) = 100^\circ$$


20.



Answer: (b) 30°

Sol:

Construction: Join OB



We know that the radius and tangent are perpendicular at the point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, In quadrilateral AOBP

$$\angle AOB + \angle OBP + \angle APB + \angle OAP = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle AOB + 90^\circ + 60^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow 240^\circ + \angle AOB = 360^\circ$$

$$\Rightarrow \angle AOB = 120^\circ$$

Now, In isosceles triangles AOB

$$\angle AOB + \angle OAB + \angle OBA = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 120^\circ + 2\angle OAB = 180^\circ \quad [\because \angle OAB = \angle OBA]$$

$$\Rightarrow \angle OAB = 30^\circ$$

21.

Answer: (c) $3\sqrt{3}$ cm

Sol:

Given, PA and PB are tangents to circle with center O and radius 3 cm and $\angle APB = 60^\circ$.

Tangents drawn from an external point are equal; so, PA = PB.

And OP is the bisector of $\angle APB$, which gives $\angle OPB = \angle OPA = 30^\circ$.

$OA \perp PA$. So, from right-angled $\triangle OPA$, we have:


$$\frac{OA}{AP} = \tan 30^\circ$$

$$\Rightarrow \frac{OA}{AP} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \frac{3}{AP} = \frac{1}{\sqrt{3}}$$

$$= AP = 3\sqrt{3} \text{ cm}$$

22.



Answer: (c) 126°

Sol:

We know that the radius and tangent are perpendicular at the point of contact

Now, In $\triangle PQA$

$$\angle PQA + \angle QAP + \angle APQ = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 90^\circ + \angle QAP + 27^\circ = 180^\circ \quad [\because \angle OAB = \angle OBA]$$


$$\Rightarrow \angle QAP = 63^\circ$$

In $\triangle PQA$ and $\triangle PRA$

$$PQ = PR \quad (\text{Tangents drawn from same external point are equal})$$

$QA = RA$ (Radio of the circle)
 $AP = AP$ (common)
 By SSS congruency
 $\Delta PQA \cong \Delta PRA$
 $\angle QAP = \angle RAP = 63^\circ$
 $\therefore \angle QAR = \angle QAP + \angle RAP = 63^\circ + 63^\circ = 126^\circ$


23.



Answer:(b)117°

Sol:

Construction: Join CA and CB



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle CAP = \angle CBP = 90^\circ$$

Since, in quadrilateral ACBP all the angles are right angles


$\therefore ACPB$ is a rectangle

Now, we know that the pair of opposite sides are equal in rectangle

$$\therefore CB = AP \text{ and } CA = BP$$

Therefore, $CB = AP = 4\text{cm}$ and $CA = BP = 4\text{cm}$

24.



Answer:(b)50°

Sol:

Given, PA and PB are two tangents to a circle with center O and $\angle APB = 80^\circ$

$$\therefore \angle APO = \frac{1}{2} \angle APB = 40^\circ$$

[Since they are equally inclined to the line segment joining the center to that point and $\angle OAP = 90^\circ$]

[Since tangents drawn from an external point are perpendicular to the radius at the point of contact]

Now, in triangle AOP:


$$\angle AOP + \angle OAP + \angle APO = 180^\circ$$

$$\Rightarrow \angle AOP + 90^\circ + 40^\circ = 180^\circ$$

$$\Rightarrow \angle AOP = 180^\circ - 130^\circ$$

$$\Rightarrow \angle AOP = 50^\circ$$

25.



Answer: (a) 32°

Sol:

We know that a chord passing through the center is the diameter of the circle.

$\therefore \angle QPR = 90^\circ$ (Angle in a semi circle is 90°)

By using alternate segment theorem

We have $\angle APQ = \angle PRQ = 58^\circ$


Now, In ΔPQR

$$\angle PQR + \angle PRQ + \angle QPR = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow \angle PQR + 58^\circ + 90^\circ = 180^\circ$$

$$\Rightarrow \angle PQR = 32^\circ$$

26.



Answer: (b) 90°

Sol:

We know that a chord passing through the center is the diameter of the circle.

$\therefore \angle DPC = 90^\circ$ (Angle in a semicircle is 90°)

Now, In $\triangle CDP$

$\angle CDP + \angle DCP + \angle DPC = 180^\circ$ [Angle sum property of a triangle]

$\Rightarrow \angle CDP + \angle DCP + 90^\circ = 180^\circ$


$\Rightarrow \angle CDP + \angle DCP = 90^\circ$

By using alternate segment theorem

We have $\angle CDP = \angle CPB$

$\therefore \angle CPB + \angle ACP = 90^\circ$

27.



Answer: (d)

Sol:

We know that a chord passing through the center is the diameter of the circle.

$\therefore \angle BAC = 90^\circ$ (Angle in a semicircle is 90°)

By using alternate segment theorem

We have $\angle PAB = \angle ACB = 67^\circ$

Now, In $\triangle ABC$

$\angle ABC + \angle ACB + \angle BAC = 180^\circ$ [Angle sum property of a triangle]

$\Rightarrow \angle ABC + 67^\circ + 90^\circ = 180^\circ$

$\Rightarrow \angle ABC = 23^\circ$

Now, $\angle BAQ = 180^\circ - \angle PAB$ [Linear pair angles]

$= 180^\circ - 67^\circ$

$= 113^\circ$


Now, In $\triangle ABQ$

$\angle ABQ + \angle AQB + \angle BAQ = 180^\circ$ [Angle sum property of a triangle]

$\Rightarrow 23^\circ + \angle AQB + 113^\circ = 180^\circ$


$\Rightarrow \angle AQB = 44^\circ$

28.



Answer: (c) 90°

Sol:



We know that tangent segments to a circle from the same external point are congruent

Therefore, we have

$$NA = NC \text{ and } NC = NB$$

We also know that angle opposite to equal sides is equal

$$\therefore \angle NAC = \angle NCA \text{ and } \angle NBC = \angle NCB$$

$$\text{Now, } \angle ANC + \angle BNC = 180^\circ \quad [\text{Linear pair angles}]$$


$$\Rightarrow \angle NBC + \angle NCB + \angle NAC + \angle NCA = 180^\circ \quad [\text{Exterior angle property}]$$

$$\Rightarrow 2\angle NCB + 2\angle NCA = 180^\circ$$

$$\Rightarrow 2(\angle NCA + \angle NCA) = 180^\circ$$

$$\Rightarrow \angle ACB = 90^\circ$$

29.



Answer: (a) 60 cm^2

Sol:

Given,

$$OQ = OR = 5 \text{ cm}, OP = 13 \text{ cm}$$

$\angle OQP = \angle ORP = 90^\circ$ (Tangents drawn from an external point are perpendicular to the radius at the point of contact)

From right – angled ΔPOQ :

$$PQ^2 = (OP^2 - OQ^2)$$

$$\Rightarrow PQ^2 = (OP^2 - OQ^2)$$

$$\Rightarrow PQ^2 = 13^2 - 5^2$$

$$\Rightarrow PQ^2 = 169 - 25$$

$$\Rightarrow PQ = 144$$

$$\Rightarrow PQ = \sqrt{144}$$

$$\Rightarrow PQ = 12 \text{ cm}$$

$$\therefore \text{ar}(\Delta OQP) = \frac{1}{2} \times PQ \times OQ$$


$$\Rightarrow \text{ar}(\Delta OQP) = \left(\frac{1}{2} \times 12 \times 5 \right) \text{cm}^2$$

$$\Rightarrow \text{ar}(\Delta OQP) = 30 \text{cm}^2$$

Similarly, $\text{ar}(\Delta ORP) = 30 \text{cm}^2$

$$\therefore \text{ar}(\text{quad. PQOR}) = (30 + 30) \text{cm}^2 = 60 \text{cm}^2$$

30.



Answer: (c) 40°

Sol:

Since, $AB \parallel PR$, BQ is transversal

$$\angle BQR = \angle ABQ = 70^\circ \quad [\text{Alternative angles}]$$

$OQ \perp PQR$ (Tangents drawn from an external point are perpendicular to the radius at the point of contact)

and $AB \parallel PQR$

$$\therefore QL \perp AB; \text{ so, } OL \perp AB$$

$$\therefore OL \text{ bisects chord } AB \quad [\text{Perpendicular drawn from the center bisects the chord}]$$

From ΔQLA and QLB :

$$\angle QLA = \angle QLB = 90^\circ$$

$$LA = LB \quad (\text{OL bisects chord AB})$$

QL is the common side.

$$\therefore \triangle QLA \cong \triangle QLB \quad [\text{By SAS congruency}]$$

$$\therefore \angle QAL = \angle QBL$$

$$\Rightarrow \angle QAB = \angle QBA$$

$\therefore \triangle AQB$ is isosceles

$$\therefore \angle LQA = \angle LQB$$

$$\angle LQP = \angle LQR = 90^\circ$$

$$\angle LQB = (90^\circ - 70^\circ) = 20^\circ$$

$$\therefore \angle LQA = \angle LQB = 20^\circ$$

$$\Rightarrow \angle LQA = \angle LQB = 20^\circ$$


$$\Rightarrow \angle AQB = \angle LQA + \angle LQB$$

$$= 40^\circ$$

31.

Answer: (d) $\sqrt{125}$ cm

Sol:



We know that the radius and tangent are perpendicular at their point of contact

In right triangle PTO

By using Pythagoras theorem, we have

$$PO^2 = OT^2 + TP^2$$


$$\Rightarrow PO^2 = 5^2 + 10^2$$

$$\Rightarrow PO^2 = 25 + 100$$

$$\Rightarrow PO^2 = 125$$

$$\Rightarrow PO = \sqrt{125} \text{ cm}$$

32.



Answer: (b) 30°

Sol:

We know that a chord passing through the center is the diameter of the circle

$\therefore \angle BPA = 90^\circ$ (Angle in a semicircle is 90°)

By using alternate segment theorem

We have $\angle APT = \angle ABP = 30^\circ$

Now, In $\triangle ABP$

$\angle PBA + \angle BPA + \angle BAP = 180^\circ$ [Angle sum property of a triangle]

$\Rightarrow 30^\circ + 90^\circ + \angle BAP = 180^\circ$


$\Rightarrow \angle BAP = 60^\circ$

Now, $\angle BAP = \angle APT + \angle PTA$

$\Rightarrow 60^\circ = 30^\circ + \angle PTA$

$\Rightarrow \angle PTA = 30^\circ$

33.



Answer: (d) 18 cm

Sol:

We know that tangent segments to a circle from the same external point are congruent.

Therefore, we have

$EK = EM = 9\text{ cm}$

Now, $EK + EM = 18\text{ cm}$

$\Rightarrow ED + DK + EF + FM = 18\text{ cm}$

$\Rightarrow ED + DH + EF + HF = 18\text{ cm}$ ($\because DK = DH$ and $FM = FH$)


$\Rightarrow ED + DF + EF = 18\text{ cm}$

\Rightarrow Perimeter of $\triangle EDF = 18\text{ cm}$

34.

Answer: (b) 135°

Sol:



Suppose PA and PB are two tangents we want to draw which inclined at an angle of 45°
 We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, in quadrilateral AOBP


$$\angle AOB + \angle OBP + \angle OAP + \angle APB = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle AOB + 90^\circ + 90^\circ + 45^\circ = 360^\circ$$

$$\Rightarrow \angle AOB + 225^\circ = 360^\circ$$

$$\Rightarrow \angle AOB = 135^\circ$$

35.



Answer:(d) 70°

Sol:

PQL is a tangent OQ is the radius; so, $\angle OQL = 90^\circ$

$$\therefore \angle OQS = (90^\circ - 50^\circ) = 40^\circ$$

Now, $OQ = OS$ (Radius of the same circle)

$$\Rightarrow \angle OSQ = \angle OQS = 40^\circ$$

Similarly, $\angle ORS = (90^\circ - 60^\circ) = 30^\circ$,

And, $OR = OS$ (Radius of the same circle)


$$\Rightarrow \angle OSR = \angle ORS = 30^\circ$$

$$\therefore \angle QSR = \angle OSQ + \angle OSR$$

$$\Rightarrow \angle QSR = (40^\circ + 30^\circ)$$


$$\Rightarrow \angle QSR = 70^\circ$$

36.



Answer: (c) 22.5 cm

Sol:



We know that tangent segments to a circle from the same external point are congruent.

Therefore, we have

$$PS = PU = x$$

$$QT = QS = 12 \text{ cm}$$

$$RT = RU = 9 \text{ cm}$$

Now,

$$Ar(\Delta PQR) = Ar(\Delta POR) + Ar(\Delta QOR) + Ar(\Delta POQ)$$

$$\Rightarrow 189 = \frac{1}{2} \times OU \times PR + \frac{1}{2} \times OT \times QR + \frac{1}{2} \times OS \times PQ$$

$$\Rightarrow 378 = 6 \times (x + 9) + 6 \times (21) + 6 \times (12 + x)$$


$$\Rightarrow 63 = x + 9 + 21 + x + 12$$

$$\Rightarrow 2x = 21$$

$$\Rightarrow x = 10.5 \text{ cm}$$

$$\text{Now, } PQ = QS + SP = 12 + 10.5 + 10.5 = 22.5 \text{ cm}$$

37.

**Answer:** (d) 7.6 cm**Sol:**


We know that tangent segments to a circle from the same external point are congruent.

Therefore, we have

$$PT = PO = 3.8 \text{ cm and } PT = PR = 3.8 \text{ cm}$$

$$\therefore QR = QP + PR = 3.8 + 3.8 = 7.6 \text{ cm}$$

38.

**Answer:** (a) 9 cm**Sol:**

Tangents drawn from an external point to a circle are equal.

$$\text{So, } AQ = AP = 5 \text{ cm}$$

$$CR = CS = 3 \text{ cm}$$

$$\text{And } BR = (BC - CR)$$

$$\Rightarrow BR = (7 - 3) \text{ cm}$$

$$\Rightarrow BR = 4 \text{ cm}$$


$$BQ = BR = 4 \text{ cm}$$

$$\therefore AB = (AQ + BQ)$$

$$\Rightarrow AB = (5 + 4) \text{ cm}$$

$$\Rightarrow AB = 9 \text{ cm}$$

39.



Answer: (c) 36 cm

Sol:

Given, $AP = 6\text{ cm}$, $BP = 5\text{ cm}$, $CQ = 3\text{ cm}$ and $DR = 4\text{ cm}$

Tangents drawn from an external point to a circle are equal

So, $AP = AS = 6\text{ cm}$, $BP = BQ = 5\text{ cm}$, $CQ = CR = 3\text{ cm}$, $DR = DS = 4\text{ cm}$.

$$\therefore AB = AP + BP = 6 + 5 = 11\text{ cm}$$

$$BC = BQ + CQ = 5 + 3 = 8\text{ cm}$$

$$CD = CR + DR = 3 + 4 = 7\text{ cm}$$


$$AD = AS + DS = 6 + 4 = 10\text{ cm}$$

$$\therefore \text{Perimeter of quadrilateral } ABCD = AB + BC + CD + DA$$

$$= (11 + 8 + 7 + 10)\text{ cm}$$

$$= 36\text{ cm}$$

40.



Answer:(b) 50°

Sol:

Given: AO and BO are the radius of the circle

Since, $AO = BO$

$\therefore \triangle AOB$ is an isosceles triangle

Now, in $\triangle AOB$

$$\angle AOB + \angle OBA + \angle OAB = 180^\circ$$

(Angle sum property of triangle)

$$\Rightarrow 100^\circ + \angle OAB + \angle OAB = 180^\circ \quad (\angle OBA = \angle OAB)$$

$$\Rightarrow 2\angle OAB = 80^\circ$$

$$\Rightarrow \angle OAB = 40^\circ$$

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OAT = 90^\circ$$


$$\Rightarrow \angle OAB + \angle BAT = 90^\circ$$

$$\Rightarrow \angle BAT = 90^\circ - 40^\circ = 50^\circ$$

41.

Answer: (b) 2 cm

Sol:



In right triangle ABC

By using Pythagoras theorem we have

$$AC^2 = AB^2 + BC^2$$

$$= 5^2 + 12^2$$

$$= 25 + 144$$

$$= 169$$

$$\therefore AC^2 = 169$$

$$\Rightarrow AC = 13 \text{ cm}$$

Now,

$$Ar(\triangle ABC) = Ar(\triangle AOB) + Ar(\triangle BOC) + Ar(\triangle AOC)$$


$$\Rightarrow \frac{1}{2} \times AB \times BC = \frac{1}{2} \times OP \times AB + \frac{1}{2} \times OQ \times BC + \frac{1}{2} \times OR \times AC$$

$$\Rightarrow 5 \times 12 = x \times 5 + x \times 12 + x \times 13$$

$$\Rightarrow 60 = 30x$$

$$\Rightarrow x = 2 \text{ cm}$$


42.



Answer: (d) 21 cm

Sol:

Construction: Join OR



We know that tangent segments to a circle from the same external point are congruent.

Therefore, we have

$$BP = BQ = 27 \text{ cm}$$

$$CQ = CR$$

Now, $BC = 38 \text{ cm}$

$$\Rightarrow BQ + QC = 38$$

$$\Rightarrow QC = 38 - 27 = 11 \text{ cm}$$

Since, all the angles in quadrilateral DROS are right angles.

Hence, DROS is a rectangle.

We know that opposite sides of rectangle are equal

$$\therefore OS = RD = 10 \text{ cm}$$


Now, $CD = CR + RD$

$$= CQ + RD$$

$$= 11 + 10$$

$$= 21 \text{ cm}$$

43.



Answer: (a) 2 cm

Sol:

Given, $AB = 8\text{ cm}$, $BC = 6\text{ cm}$

Now, in $\triangle ABC$:

$$AC^2 = AB^2 + BC^2$$

$$\Rightarrow AC^2 = (8^2 + 6^2)$$

$$\Rightarrow AC^2 = (64 + 36)$$

$$\Rightarrow AC^2 = 100$$

$$\Rightarrow AC = \sqrt{100}$$

$$\Rightarrow AC = 10\text{ cm}$$

$PBQO$ is a square

$CR = CQ$ (Since the lengths of tangents drawn from an external point are equal)

$$\therefore CQ = (BC - BQ) = (6 - x)\text{ cm}$$

Similarly, $AR = AP = (AB - BP) = (8 - x)\text{ cm}$

$$\therefore AC = (AR + CR) = [(8 - x) + (6 - x)]\text{ cm}$$

$$\Rightarrow 10 = (14 - 2x)\text{ cm}$$

$$\Rightarrow 2x = 4$$


$$\Rightarrow x = 2\text{ cm}$$

\therefore The radius of the circle is 2 cm.

44.

Answer: (a) 3 cm

Sol:




We know that when a quadrilateral circumscribes a circle then sum of opposite sides is equal to the sum of other opposite sides

$$\therefore AB + DC = AD + BC$$

$$\Rightarrow 6 + 4 = AD + 7$$

$$\Rightarrow AD = 3\text{ cm}$$

45.

**Answer:** (b) 5 cm**Sol:**

The lengths of tangents drawn from a point to a circle are equal
So, $PA = PB$ and therefore, $\angle PAB = \angle PBA = x$ (say).

Then, in $\triangle PAB$:

$$\angle PAB + \angle PBA + \angle APB = 180^\circ$$

$$\Rightarrow x + x + 60^\circ = 180^\circ$$

$$\Rightarrow 2x = 180^\circ - 60^\circ$$

$$\Rightarrow 2x = 120^\circ$$

$$\Rightarrow x = 60^\circ$$

\therefore Each angle of $\triangle PAB$ is 60° and therefore, it is an equilateral triangle.


$\therefore AB = PA = PB = 5 \text{ cm}$

\therefore The length of the chord AB is 5 cm .

46.

Answer: (c) 5 cm**Sol:**

Construction: Join AF and AE



We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle AED = \angle AFD = 90^\circ$$

Since, in quadrilateral $AEDF$ all the angles are right angles


$\therefore AEDF$ is a rectangle

Now, we know that the pair of opposite sides is equal in rectangle

$$\therefore AF = DE = 5 \text{ cm}$$

Therefore, the radius of the circle is 5 cm

47.



Answer: (b) 2 cm

Sol:

Given, $AB = 5$ cm, $BC = 7$ cm and $CA = 6$ cm.

Let, $AR = AP = x$ cm.

$BQ = BP = y$ cm

$CR = CQ = z$ cm

(Since the length of tangents drawn from an external point are equal)

Then, $AB = 5$ cm

$\Rightarrow AP + PB = 5$ cm

$\Rightarrow x + y = 5$ (i)

Similarly, $y + z = 7$ (ii)

and $z + x = 6$ (iii)

Adding (i), (ii) and (iii), we get:

$$(x + y) + (y + z) + (z + x) = 18$$

$$\Rightarrow 2(x + y + z) = 18$$


$$\Rightarrow (x + y + z) = 9$$
(iv)

Now, (iv) - (ii):

$$\Rightarrow x = 2$$


\therefore The radius of the circle with center A is 2 cm.

48.



Answer: (d) 7.5 cm

Sol:



We know that tangent segments to a circle from the same external point are congruent

Therefore, we have

$$AP = AQ$$

$$BP = BD$$

$$CQ = CD$$

$$\text{Now, } AB + BC + AC = 5 + 4 + 6 = 15$$

$$\Rightarrow AB + BD + DC + AC = 15 \text{ cm}$$


$$\Rightarrow AB + BP + CQ + AC = 15 \text{ cm}$$

$$\Rightarrow AP + AQ = 15 \text{ cm}$$

$$\Rightarrow 2AP = 15 \text{ cm}$$


$$\Rightarrow AP = 7.5 \text{ cm}$$

49.



Answer: (c) $4\sqrt{10}$ cm

Sol:



Given, $OP = 5$ cm, $PA = 12$ cm

Now, join O and B

Then, $OB = 3 \text{ cm}$.

Now, $\angle OAP = 90^\circ$ (Tangents drawn from an external point are perpendicular to the radius at the point of contact)

Now, in $\triangle OAP$:

$$OP^2 = OA^2 + PA^2$$

$$\Rightarrow OP^2 = 5^2 + 12^2$$

$$\Rightarrow OP^2 = 25 + 144$$

$$\Rightarrow OP^2 = 169$$

$$\Rightarrow OP = \sqrt{169}$$

$$\Rightarrow OP = 13$$

Now, in $\triangle OBP$:

$$PB^2 = OP^2 - OB^2$$

$$\Rightarrow PB^2 = 13^2 - 3^2$$

$$\Rightarrow PB^2 = 169 - 9$$

$$\Rightarrow PB^2 = 160$$

$$\Rightarrow PB = \sqrt{160}$$

$$\Rightarrow PB = 4\sqrt{10} \text{ cm}$$

50.

Answer: (d) A circle can have more than two parallel tangents. parallel to a given line.

Sol:

A circle can have more than two parallel tangents. parallel to a given line.

This statement is false because there can only be two parallel tangents to the given line in a circle.

51.

Answer: (d) A straight line can meet a circle at one point only.

Sol:

A straight line can meet a circle at one point only

This statement is not true because a straight line that is not a tangent but a secant cuts the circle at two points.

52.

Answer: (d) A tangent to the circle can be drawn from a point inside the circle.

Sol:


A tangent to the circle can be drawn from a point Inside the circle.

This statement is false because tangents are the lines drawn from an external point to the circle that touch the circle at one point.

53.

Answer: (a) Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A).

Sol:



(a) Both Assertion (A) and Reason (R) are true and Reason (R) is a correct explanation of Assertion (A)

In $\triangle OPQ$, $\angle OPQ = 90^\circ$

$$\begin{aligned} \therefore OQ^2 &= OP^2 + PQ^2 \\ \Rightarrow OQ &= \sqrt{OP^2 + PQ^2} \\ &= \sqrt{12^2 + 16^2} \\ &= \sqrt{144 + 256} \\ &= \sqrt{400} \\ &= 20 \text{ cm} \end{aligned}$$

54.


Answer: (b) Both Assertion (A) and Reason (R) are true but Reason (R) is not a correct explanation of Assertion (A).

Sol:

Assertion -

We know that If two tangents are drawn to a circle from an external point, they subtend equal angles at the center

Reason:



Given, a parallelogram $ABCD$ circumscribes a circle with center O

$$AB = BC = CD = AD$$

We know that the tangents drawn from an external point to circle are equal

$$\therefore AP = AS \quad \dots\dots\dots(i) \quad \text{[tangents from } A\text{]}$$

$$BP = BQ \quad \dots\dots\dots(ii) \quad \text{[tangents from } B\text{]}$$

$$CR = CQ \quad \dots\dots\dots(iii) \quad \text{[tangents from } C\text{]}$$

$$DR = DS \quad \dots\dots\dots(iv) \quad \text{[tangents from } D\text{]}$$

$$\begin{aligned} \therefore AB + CD &= AP + BP + CR + DR \\ &= AS + BQ + CQ + DS \end{aligned} \quad \text{[from (i), (ii), (iii) and (iv)]}$$

$$= (AS + DS) + (BQ + CQ)$$

$$= AD + BC$$


55.

The correct answer is (a) / (b) / (c) / (d).

Answer: (d) Assertion (A) is false and Reason (R) is true.

Exercise - Formative Assessment

1.



Answer: (b) 100°

Sol:

Given, $\angle QPT = 50^\circ$

Now, $\angle OPT = 90^\circ$ (Since tangents drawn from an external point are perpendicular to the radius at point of contact)

$$\therefore \angle OPQ = (\angle OPT - \angle QPT) = (90^\circ - 50^\circ) = 40^\circ$$

$$OP = OQ \quad (\text{Radii of the same circle})$$

$$\Rightarrow \angle OPQ = \angle OQP = 40^\circ$$

In $\triangle POQ$

$$\angle POQ + \angle OPQ + \angle OQP = 180^\circ$$

$$\Rightarrow \angle POQ + 40^\circ + 40^\circ = 180^\circ$$

$$\Rightarrow \angle POQ = 180^\circ - (40^\circ + 40^\circ)$$


$$\Rightarrow \angle POQ = 180^\circ - 80^\circ$$

$$\Rightarrow \angle POQ = 100^\circ$$

2.

Answer: (c) 50°

Sol:



OA and OB are the two radii of a circle with center O .

Also, AP and BP are the tangents to the circle.

Given, $\angle AOB = 130^\circ$

Now, $\angle OAB = \angle OBA = 90^\circ$ (Since tangents drawn from an external point are perpendicular to the radius at point of contact)

In quadrilateral $OAPB$,

$$\angle AOB + \angle OAB + \angle OBA + \angle APB = 360^\circ$$


$$\Rightarrow 130^\circ + 90^\circ + 90^\circ + \angle APB = 360^\circ$$

$$\Rightarrow \angle APB = 360^\circ - (130^\circ + 90^\circ + 90^\circ)$$

$$\Rightarrow \angle APB = 360^\circ - 310^\circ$$

$$\Rightarrow \angle APB = 50^\circ$$

3.



Answer: (b) 50°

Sol:From $\triangle OPA$ and $\triangle OPB$

$$OA = OB \quad (\text{Radii of the same circle})$$

$$OP \quad (\text{Common side})$$

$$PA = PB \quad (\text{Since tangents drawn from an external point to a circle are equal})$$

$$\therefore \triangle OPA \cong \triangle OPB \quad (\text{SSS rule})$$

$$\therefore \angle APO = \angle BPO$$

$$\therefore \angle APO = \frac{1}{2} \angle APB = 40^\circ$$


And $\angle OAP = 90^\circ$ (Since tangents drawn from an external point are perpendicular to the radius at point of contact)

$$\text{Now, in } \triangle OAP, \angle AOP + \angle OAP + \angle APO = 180^\circ$$

$$\Rightarrow \angle AOP + 90^\circ + 40^\circ = 180^\circ$$

$$\Rightarrow \angle AOP = 180^\circ - 130^\circ = 50^\circ$$

4.

**Answer:** (b) 10cm**Sol:**

Since the tangents from an external point are equal, we have

$$AD = AE, CD = CF, BE = BF$$

$$\text{Perimeter of } \triangle ABC = AC + AB + CB$$

$$= (AD - CD) + (CF + BF) + (AE - BE)$$

$$= (AD - CF) + (CF + BF) + (AE - BF)$$


$$= AD + AE$$

$$= 2AE$$

$$= 2 \times 5$$

$$= 10 \text{ cm}$$

5.



Sol:

We know that tangent segments to a circle from the same external point are congruent

Now, we have

$$CR = CQ, AS = AP \text{ and } BQ = BP$$

Now, $BC = 7 \text{ cm}$

$$\Rightarrow CQ + BQ = 7$$

$$\Rightarrow BQ = 7 - CQ$$

$$\Rightarrow BQ = 7 - 3 \quad [\because CQ = CR = 3]$$

$$\Rightarrow BQ = 4 \text{ cm}$$

Again, $AB = AP + PB$


$$= AP = BQ$$

$$= 5 + 4 \quad [\because AS = AP = 5]$$

$$= 9 \text{ cm}$$

Hence, the value of x is 9 cm

6.



Sol:

Here, $OA = OB$

And $OA \perp AP, OA \perp BP$, (Since tangents drawn from an external point are perpendicular to the radius at the point of contact)

$$\therefore \angle OAP = 90^\circ, \angle OBP = 90^\circ$$


$$\therefore \angle OAP + \angle OBP = 90^\circ + 90^\circ = 180^\circ$$

$$\therefore \angle AOB + \angle APB = 180^\circ \quad (\text{Since, } \angle OAP + \angle OBP + \angle AOB + \angle APB = 360^\circ)$$

Sum of opposite angle of a quadrilateral is 180° .

Hence, A, O, B and P are concyclic.

7.



Sol:

We know that tangents drawn from the external point are congruent

$$\therefore PA = PB$$

Now, In isosceles triangle APB

$$\angle APB + \angle PBA + \angle PAB = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow \angle APB + 65^\circ + 65^\circ = 180^\circ \quad [\because \angle PBA = \angle PAB = 65^\circ]$$

$$\Rightarrow \angle APB = 50^\circ$$

We know that the radius and tangent are perpendicular at their point of contact

$$\therefore \angle OBP = \angle OAP = 90^\circ$$

Now, In quadrilateral AOBP

$$\angle AOB + \angle OBP + \angle APB + \angle OAP = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow \angle AOB + 90^\circ + 50^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow 230^\circ + \angle BOC = 360^\circ$$

$$\Rightarrow \angle AOB = 130^\circ$$


Now, In isosceles triangle AOB

$$\angle AOB + \angle OAB + \angle OBA = 180^\circ \quad [\text{Angle sum property of a triangle}]$$

$$\Rightarrow 130^\circ + 2\angle OAB = 180^\circ \quad [\because \angle OAB = \angle OBA]$$

$$\Rightarrow \angle OAB = 25^\circ$$

8.



Ans:

Sol:

Here, OB is the bisector of $\angle CBD$.

(Two tangents are equally inclined to the line segment joining the center to that point)

$$\therefore \angle CBO = \angle DBO = \frac{1}{2} \angle CBD = 60^\circ$$

$$\therefore \text{From } \triangle BOD, \angle BOD = 30^\circ$$

Now, from right – angled $\triangle BOD$,

$$\Rightarrow \frac{BD}{OB} = \sin 30^\circ$$

$$\Rightarrow OB = 2BD$$

$$\Rightarrow OB = 2BC \text{ (Since tangents from an external point are equal. i.e., } BC = BD)$$

$$\therefore OB = 2BC$$

9.

Sol:

(i) A line intersecting a circle at two distinct points is called a secant


(ii) A circle can have two parallel tangents at the most

(iii) The common point of a tangent to a circle and the circle is called the point of contact.

(iv) A circle can have infinite tangents

10.

Sol:



Given two tangents AP and AQ are drawn from a point A to a circle with center O.

To prove: $AP = AQ$

Join OP , OQ and OA .

AP is tangent at P and OP is the radius.

$\therefore OP \perp AP$ (Since tangents drawn from an external point are perpendicular to the radius at the point of contact)

Similarly, $OQ \perp AQ$

In the right $\triangle OPA$ and $\triangle OQA$, we have:

$$OP = OQ \quad \text{[radii of the same circle]}$$

$$\angle OPA = \angle OQA (= 90^\circ)$$


$$OA = OA \quad \text{[Common side]}$$

$$\therefore \triangle OPA \cong \triangle OQA \quad \text{[By R.H.S – Congruence]}$$

Hence, $AP = AQ$

11.

Sol:



Here, PT and QS are the tangents to the circle with center O and AB is the diameter

Now, radius of a circle is perpendicular to the tangent at the point of contact


$\therefore OA \perp AT$ and $OB \perp BS$ (Since tangents drawn from an external point are perpendicular to the radius at point of contact)

$\therefore \angle OAT = \angle OBQ = 90^\circ$

But $\angle OAT$ and $\angle OBQ$ are alternate angles.

$\therefore AT$ is parallel to BS .

12.



Sol:

Given, $AB = AC$

We know that the tangents from an external point are equal

$\therefore AD = AF, BD = BE$ and $CF = CE$ (i)

Now, $AB = AC$

$\Rightarrow AD + DB = AF + FC$

$\Rightarrow AF + DB = AF + FC$ [from(i)]


$\Rightarrow DB = FC$

$\Rightarrow BE = CE$ [from(i)]

Hence proved.

13.

Sol:



Given: A circle with center O and a point A outside it. Also, AP and AQ are the two tangents to the circle

To prove: $\angle AOP = \angle AOQ$.

Proof : In $\triangle AOP$ and $\triangle AOQ$, we have

$AP = AQ$ [tangents from an external point are equal]

$OP = OQ$ [radii of the same circle]


$OA = OA$ [common side]

$\therefore \triangle AOP \cong \triangle AOQ$ [by SSS – congruence]

Hence, $\angle AOP = \angle AOQ$ (c.p.c.t).

14.

Sol:



Let RA and RB be two tangents to the circle with center O and let AB be a chord of the circle.

We have to prove that $\angle RAB = \angle RDA$.


\therefore Now, RA

$= RB$ (Since tangents drawn from an external point to a circle are equal)

In $\triangle RAB$, $\angle RAB = \angle RDA$ (Since opposite sides are equal, their base angles are also equal)

15.

Sol:



Given, a parallelogram $ABCD$ circumscribes a circle with center O

$$AB = BC = CD = AD$$

We know that the tangents drawn from an external point to circle are equal

$$\therefore AP = AS \quad \dots\dots\dots(i) \quad \text{[tangents from } A\text{]}$$

$$BP = BQ \quad \dots\dots\dots(ii) \quad \text{[tangents from } B\text{]}$$

$$CR = CQ \quad \dots\dots\dots(iii) \quad \text{[tangents from } C\text{]}$$

$$DR = DS \quad \dots\dots\dots(iv) \quad \text{[tangents from } D\text{]}$$

$$\begin{aligned} \therefore AB + CD &= AP + BP + CR + DR \\ &= AS + BQ + CQ + DS \end{aligned} \quad \text{[from (i), (ii), (iii) and (iv)]}$$

$$= (AS + DS) + (BQ + CQ)$$

$$= AD + BC$$

$$\text{Thus, } (AB + CD) = (AD + BC)$$

$$\Rightarrow 2AB = 2AD \quad \text{[}\because \text{ opposite sides of a parallelogram are equal]}$$


$$\Rightarrow AB = AD$$

$$\therefore CD = AB = AD = BC$$

Hence, $ABCD$ is a rhombus.

16.

Sol:



Given: Two circles have the same center O and AB is a chord of the larger circle touching the smaller circle at C . also, $OA = 5$ cm and $OC = 3$ cm

$$\text{In } \triangle OAC, OA^2 = OC^2 + AC^2$$

$$\therefore AC^2 = OA^2 - OC^2$$

$$\Rightarrow AC^2 = 5^2 - 3^2$$

$$\Rightarrow AC^2 = 25 - 9$$

$$\Rightarrow AC^2 = 16$$

$$\Rightarrow AC = 4 \text{ cm}$$


$$\therefore AB = 2AC \text{ (Since perpendicular drawn from the center of the circle bisects the chord)}$$

$$\therefore AB = 2 \times 4 = 8 \text{ cm}$$

The length of the chord of the larger circle is 8cm.

17.

Sol:



We know that the tangents drawn from an external point to circle are equal.

$$\therefore AP = AS \quad \dots\dots\dots(i) \quad \text{[tangents from } A\text{]}$$

$$BP = BQ \quad \dots\dots\dots(ii) \quad \text{[tangents from } B\text{]}$$

$$CR = CQ \quad \dots\dots\dots(iii) \quad \text{[tangents from } C\text{]}$$


$$DR = DS \quad \dots\dots\dots(iv) \quad \text{[tangents from } D\text{]}$$

$$\begin{aligned} \therefore AB + CD &= (AP + BP) + (CR + DR) \\ &= (AS + BQ) + (CQ + DS) \quad \text{[using (i), (ii), (iii) and (iv)]} \\ &= (AS + DS) + (BQ + CQ) \\ &= AD + BC \end{aligned}$$

Hence, $(AB + CD) = (AD + BC)$

18.

Sol:



Given, a quadrilateral $ABCD$ circumscribes a circle with center O .

To prove: $\angle AOB + \angle COD = 180^\circ$

And $\angle AOD + \angle BOC = 180^\circ$

Join: OP, OQ, OR and OS .

We know that the tangents drawn from an external point of a circle subtend equal angles at the center.


$$\therefore \angle 1 = \angle 7, \angle 2 = \angle 3, \angle 4 = \angle 5 \text{ and } \angle 6 = \angle 8$$

And $\angle 1 + \angle 2 + \angle 3 + \angle 4 + \angle 5 + \angle 6 + \angle 7 + \angle 8 = 360^\circ$ [angles at a point]
 $\Rightarrow (\angle 1 + \angle 7) + (\angle 3 + \angle 2) + (\angle 4 + \angle 5) + (\angle 6 + \angle 8) = 360^\circ$
 $2\angle 1 + 2\angle 2 + 2\angle 6 + 2\angle 5 = 360^\circ$
 $\Rightarrow \angle 1 + \angle 2 + \angle 5 + \angle 6 = 180^\circ$
 $\Rightarrow \angle AOB + \angle COD = 180^\circ$ and $\angle AOD + \angle BOC = 180^\circ$

19.

Ans:

Sol:



Given, PA and PB are the tangents drawn from a point P to a circle with center O . Also, the line segments OA and OB are drawn.

To prove: $\angle APB + \angle AOB = 180^\circ$

We know that the tangent to a circle is perpendicular to the radius through the point of contact

$\therefore PA \perp OA$

$\Rightarrow \angle OAP = 90^\circ$

$PB \perp OB$

$\Rightarrow \angle OBP = 90^\circ$

$\therefore \angle OAP + \angle OBP = (90^\circ + 90^\circ) = 180^\circ$ (i)


But we know that the sum of all the angles of a quadrilateral is 360° .

$\therefore \angle OAP + \angle OBP + \angle APB + \angle AOB = 360^\circ$ (ii)


From (i) and (ii), we get:

$\angle APB + \angle AOB = 180^\circ$

20.



Sol:



Let $TR = y$ and $TP = x$

We know that the perpendicular drawn from the center to the chord bisects it.

$\therefore PR = RQ$

Now, $PR + RQ = 16$

$PR + PR = 16$

$\Rightarrow PR = 8$

Now, in right triangle POR

By Using Pythagoras theorem, we have

$$PO^2 = OR^2 + PR^2$$

$$\Rightarrow 10^2 = OR^2 + (8)^2$$

$$\Rightarrow OR^2 = 36$$

$$\Rightarrow OR = 6$$

Now, in right triangle TPR

By Using Pythagoras theorem, we have

$$TP^2 = TR^2 + PR^2$$

$$\Rightarrow x^2 = y^2 + (8)^2$$

$$\Rightarrow x^2 = y^2 + 64 \quad \dots\dots(1)$$

Again, in right triangle TPQ

By Using Pythagoras theorem, we have

$$TO^2 = TP^2 + PO^2$$

$$\Rightarrow (y+6)^2 = x^2 + 10^2$$

$$\Rightarrow y^2 + 12y + 36 = x^2 + 100$$

$$\Rightarrow y^2 + 12y = x^2 + 64 \quad \dots\dots(2)$$

Solving (1) and (2), we get

$$x = 10.67$$

$\therefore TP = 10.67 \text{ cm}$