

## Exercise 9A

1. **Sol:**

$$\text{Mean of given observations} = \frac{\text{sum of given observations}}{\text{total number of observations}}$$

$$\therefore 11 = \frac{x + (x+2) + (x+4) + (x+6) + (x+8)}{5}$$

$$\Rightarrow 55 = 5x + 20$$

$$\Rightarrow 5x = 55 - 20$$

$$\Rightarrow 5x = 35$$

$$\Rightarrow x = \frac{35}{5}$$

$$\Rightarrow x = 7$$

Hence, the value of  $x$  is 7.

2.

**Sol:**

$$\text{Mean of given observations} = \frac{\text{sum of given observations}}{\text{total number of observations}}$$

$$\text{Mean of 25 observations} = 27$$

$$\therefore \text{Sum of 25 observations} = 27 \times 25 = 675$$

$$\begin{aligned} \text{If 7 is subtracted from every number, then the sum} &= 675 - (25 \times 7) \\ &= 675 - 175 \\ &= 500 \end{aligned}$$

$$\text{Then, new mean} = \frac{500}{25} = 20$$

Thus, the new mean will be 20.

3.

**Sol:**

The given data is shown as follows:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
1 - 3	12	2	24
3 - 5	22	4	88
5 - 7	27	6	162
7 - 9	19	8	152
Total	$\Sigma f_i = 80$		$\Sigma f_i x_i = 426$

The mean of given data is given by

$$\begin{aligned}\bar{x} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ &= \frac{426}{80} \\ &= 5.325\end{aligned}$$

Thus, the mean of the following data is 5.325.

4.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$f_i \times x_i$
0 – 10	7	5	35
10 – 20	5	15	75
20 – 30	6	25	150
30 – 40	12	35	420
40 – 50	8	45	360
50 – 60	2	55	110
	$\Sigma f_i = 40$		$\Sigma(f_i \times x_i) = 1150$

$$\therefore \text{Mean, } \bar{x} = \frac{\Sigma(f_i \times x_i)}{\Sigma f_i}$$

$$= \frac{1150}{40}$$

$$= 28.75$$

$$\therefore \bar{x} = 28.75$$

5.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$(f_i \times x_i)$
25 – 35	6	30	180
35 – 45	10	40	400
45 – 55	8	50	400
55 – 65	12	60	720
65 – 75	4	70	280
	$\Sigma f_i = 40$		$\Sigma(f_i \times x_i) = 1980$

$$\therefore \text{Mean, } \bar{x} = \frac{\sum(f_i \times x_i)}{\sum f_i}$$

$$= \frac{1980}{40}$$

$$= 49.5$$

$$\therefore \bar{x} = 49.5$$

6.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	( $f_i \times x_i$ )
0 - 100	6	50	300
100 - 200	9	150	1350
200 - 300	15	250	3750
300 - 400	12	350	4200
400 - 500	8	450	3600
	$\Sigma f_i = 50$		$\Sigma(f_i \times x_i) = 13200$

$$\therefore \text{Mean, } \bar{x} = \frac{\sum(f_i \times x_i)}{\sum f_i}$$

$$= \frac{13200}{50}$$

$$= 264$$

$$\therefore \bar{x} = 264$$

7.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	( $f_i \times x_i$ )
84 - 90	8	87	696
90 - 96	10	93	930
96 - 102	16	99	1584
102 - 108	23	105	2415
108 - 114	12	111	1332
114 - 120	11	117	1287
Total	$\Sigma f_i = 80$		$\Sigma f_i \times x_i = 8244$

The mean of the data is given by,

$$\begin{aligned}\bar{x} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ &= \frac{8244}{80} \\ &= 103.05\end{aligned}$$

Thus, the mean of the following data is 103.05.

8.

**Sol:**

The given data is shown as follows:

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	( $f_i x_i$ )
0 – 10	3	5	15
10 – 20	4	15	60
20 – 30	p	25	25p
30 – 40	3	35	105
40 – 50	2	45	90
Total	$\sum f_i = 12 + p$		$\sum f_i x_i = 270 + 25p$

The mean of the given data is given by,

$$\begin{aligned}\bar{x} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ \Rightarrow 24 &= \frac{270+25p}{12+p}\end{aligned}$$

$$\Rightarrow 24(12 + p) = 270 + 25p$$

$$\Rightarrow 288 + 24p = 270 + 25p$$

$$\Rightarrow 25p - 24p = 288 - 270$$

$$\Rightarrow p = 18$$

Hence, the value of p is 18.

9.

**Sol:**

The given data is shown as follows:

Daily pocket allowance (in ₹)	Number of children ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
11 – 13	7	12	84
13 – 15	6	14	84
15 – 17	9	16	144
17 – 19	13	18	234
19 – 21	$f$	20	$20f$
21 – 23	5	22	110
23 – 25	4	24	96
Total	$\Sigma f_i = 44 + f$		$\Sigma f_i x_i = 752 + 20f$

The mean of the given data is given by,

$$\bar{x} = \frac{\sum_i f_i x_i}{\sum_i f_i}$$

$$\Rightarrow 18 = \frac{750+20f}{44+f}$$

$$\Rightarrow 18(44 + f) = 752 + 20f$$

$$\Rightarrow 792 + 18f = 752 + 20f$$

$$\Rightarrow 20f - 18f = 792 - 752$$

$$\Rightarrow 2f = 40$$

$$\Rightarrow f = 20$$

Hence, the value of  $f$  is 20.

10.

**Sol:**

The given data is shown as follows:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
0 – 20	7	10	70
20 – 40	$p$	30	$30p$
40 – 60	10	50	500
60 – 80	9	70	630
80 – 100	13	90	1170
Total	$\Sigma f_i = 39 + p$		$\Sigma f_i x_i = 2370 + 30p$

The mean of the given data is given by,

$$\bar{x} = \frac{\sum_i f_i x_i}{\sum_i f_i}$$

$$\Rightarrow 54 = \frac{2370 + 30p}{39 + p}$$

$$\Rightarrow 54(39 + p) = 2370 + 30p$$

$$\Rightarrow 2106 + 54p = 2370 - 2106$$

$$\Rightarrow 24p = 264$$

$$\Rightarrow p = 11$$

Hence, the value of p is 11.

11.

**Sol:**

The given data is shown as follows:

Class interval	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
0 – 10	7	5	35
10 – 20	10	15	150
20 – 30	x	25	25x
30 – 40	13	35	455
40 – 50	y	45	45y
50 – 60	10	55	550
60 – 70	14	65	910
70 – 80	9	75	675
Total	$\sum f_i = 63 + x + y$		$\sum f_i x_i = 2775 + 25x + 45y$

Sum of the frequencies = 100

$$\Rightarrow \sum_i f_i = 100$$

$$\Rightarrow 63 + x + y = 100$$

$$\Rightarrow x + y = 100 - 63$$

$$\Rightarrow x + y = 37$$

$$\Rightarrow y = 37 - x \quad \dots\dots\dots(1)$$

Now, the mean of the given data is given by,

$$\bar{x} = \frac{\sum_i f_i x_i}{\sum_i f_i}$$

$$\Rightarrow 42 = \frac{2775 + 25x + 45y}{100}$$

$$\Rightarrow 4200 = 2775 + 25x + 45y$$

$$\Rightarrow 4200 - 2775 = 25x + 45y$$

$$\Rightarrow 1425 = 25x + 45(37 - x) \quad [\text{from (1)}]$$

$$\Rightarrow 1425 = 25x + 1665 - 45x$$

$$\Rightarrow 20x = 1665 - 1425$$

$$\Rightarrow 20x = 240$$

$$\Rightarrow x = 12$$

$$\text{If } x = 12, \text{ then } y = 37 - 12 = 25$$

Thus, the value of x is 12 and y is 25.

12.

**Sol:**

The given data is shown as follows:

Expenditure (in ₹)	Number of families ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
140 – 160	5	150	750
160 – 180	25	170	4250
180 – 200	$f_1$	190	$190f_1$
200 – 220	$f_2$	210	$210f_2$
220 – 240	5	230	1150
Total	$\Sigma f_i = 35 + f_1 + f_2$		$\Sigma f_i x_i = 6150 + 190f_1 + 210f_2$

Sum of the frequencies = 100

$$\Rightarrow \Sigma_i f_i = 100$$

$$\Rightarrow 35 + f_1 + f_2 = 100$$

$$\Rightarrow f_1 + f_2 = 100 - 35$$

$$\Rightarrow f_1 + f_2 = 65$$

$$\Rightarrow f_2 = 65 - f_1 \quad \dots\dots(1)$$

Now, the mean of the given data is given by,

$$\bar{x} = \frac{\sum_i f_i x_i}{\sum_i f_i}$$

$$\Rightarrow 188 = \frac{6150 + 190f_1 + 210f_2}{100}$$

$$\Rightarrow 18800 = 6150 + 190f_1 + 210f_2$$

$$\Rightarrow 18800 - 6150 = 190f_1 + 210f_2$$

$$\Rightarrow 12650 = 190f_1 + 210(65 - f_1) \quad [\text{from (1)}]$$

$$\Rightarrow 12650 = 190f_1 - 210f_1 + 13650$$

$$\Rightarrow 20f_1 = 13650 - 12650$$

$$\Rightarrow 20f_1 = 1000$$

$$\Rightarrow f_1 = 50$$

If  $f_1 = 50$ , then  $f_2 = 65 - 50 = 15$

Thus, the value of  $f_1$  is 50 and  $f_2$  is 15.

13.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	( $f_i \times x_i$ )
0 - 20	7	10	70
20 - 40	$f_1$	30	$30f_1$
40 - 60	12	50	600
60 - 80	$18 - f_1$	70	$1260 - 70f_1$
80 - 100	8	90	720
100 - 120	5	110	550
Total	$\Sigma f_i = 50$		$\Sigma (f_i \times x_i) = 3200 - 40f_1$

We have:

$$7 + f_1 + 12 + f_2 + 8 + 5 = 50$$

$$\Rightarrow f_1 + f_2 = 18$$

$$\Rightarrow f_2 = 18 - f_1$$

$$\therefore \text{Mean, } \bar{x} = \frac{\Sigma_i (f_i \times x_i)}{\Sigma_i f_i}$$

$$\Rightarrow 57.6 = \frac{3200 - 40f_1}{50}$$



$$\Rightarrow 40f_1 = 320$$

$$\therefore f_1 = 8$$

$$\text{And } f_2 = 18 - 8$$

$$\Rightarrow f_2 = 10$$

$\therefore$  The missing frequencies are  $f_1 = 8$  and  $f_2 = 10$ .

14.

**Sol:**

Using Direct method, the given data is shown as follows:

Number of heartbeats per minute	Number of patients ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
65 – 68	2	66.5	133
68 – 71	4	69.5	278
71 – 74	3	72.5	217.5
74 – 77	8	75.5	604
77 – 80	7	78.5	549.5
80 – 83	4	81.5	326
83 – 86	2	84.5	169
Total	$\Sigma f_i = 30$		$\Sigma f_i x_i = 2277$

The mean of the data is given by,

$$\begin{aligned}\bar{x} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ &= \frac{2277}{30} \\ &= 75.9\end{aligned}$$

Thus, the mean heartbeats per minute for these patients is 75.9.

15.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	Deviation ( $d_i$ ) $d_i = (x_i - 25)$	$(f_i \times d_i)$
0 – 10	12	5	-20	-240
10 – 20	18	15	-10	-180
20 – 30	27	25 = A	0	0
30 – 40	20	35	10	200
40 – 50	17	45	20	340
50 – 60	6	55	30	180
Total	$\Sigma f_i = 100$			$\Sigma (f_i \times d_i) = 300$

Let A = 25 be the assumed mean. Then we have:

$$\begin{aligned} \text{Mean, } \bar{x} &= A + \frac{\sum (f_i \times d_i)}{\sum f_i} \\ &= 25 + \frac{300}{100} \\ &= 28 \\ \therefore \bar{x} &= 28 \end{aligned}$$

16.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	Deviation ( $d_i$ ) $d_i = (x_i - 150)$	$(f_i \times d_i)$
100 – 120	10	110	-40	-400
120 – 140	20	130	-20	-400
140 – 160	30	150 = A	0	0
160 – 180	15	170	20	300
180 – 200	5	190	40	200
	$\Sigma f_i = 80$			$\Sigma (f_i \times d_i) = -300$

Let A = 150 be the assumed mean. Then we have:

$$\begin{aligned} \text{Mean, } \bar{x} &= A + \frac{\sum (f_i \times d_i)}{\sum f_i} \\ &= 150 - \frac{300}{80} \\ &= 150 - 3.75 \\ \therefore \bar{x} &= 146.25 \end{aligned}$$

17.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	Deviation ( $d_i$ ) $d_i = (x_i - 50)$	$(f_i \times d_i)$
0 – 20	20	10	-40	-800
20 – 40	35	30	-20	-700
40 – 60	52	50 = A	0	0
60 – 80	44	70	20	880
80 – 100	38	90	40	1520
100 – 120	31	110	60	1860
	$\Sigma f_i = 220$			$\Sigma (f_i \times d_i) = 2760$

Let A = 50 be the assumed mean. Then we have:

$$\begin{aligned} \text{Mean, } \bar{x} &= A + \frac{\sum (f_i \times d_i)}{\sum f_i} \\ &= 50 + \frac{2760}{220} \\ &= 50 + 12.55 \\ \therefore \bar{x} &= 62.55 \end{aligned}$$

18.

**Sol:**

Using Direct method, the given data is shown as follows:

Literacy rate (%)	Number of cities ( $f_i$ )	Class mark ( $x_i$ )	$(f_i \times x_i)$
45 – 55	4	50	200
55 – 65	11	60	660
65 – 75	12	70	840
75 – 85	9	80	720
85 – 95	4	90	360
Total	$\Sigma f_i = 40$		$\Sigma f_i x_i = 2780$

The mean of the data is given by,

$$\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$$

$$= \frac{2780}{40}$$

$$= 69.5$$

Thus, the mean literacy rate is 69.5%.

19.

**Sol:**

Let us choose  $a = 25$ ,  $h = 10$ , then  $d_i = x_i - 25$  and  $u_i = \frac{x_i - 25}{10}$

Using step-deviation method, the given data is shown as follows:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 25$	$u_i = \frac{x_i - 25}{10}$	$(f_i u_i)$
0 – 10	7	5	-20	-2	-14
10 – 20	10	15	-10	-1	-10
20 – 30	15	25	0	0	0
30 – 40	8	35	10	1	8
40 – 50	10	45	20	2	20
Total	$\Sigma f_i = 50$				$\Sigma f_i u_i = 4$

The mean of the data is given by,

$$\bar{x} = a + \left( \frac{\Sigma f_i u_i}{\Sigma f_i} \right) \times h$$

$$= 25 + \frac{4}{50} \times 10$$

$$= 25 + \frac{4}{5}$$

$$= \frac{125 + 4}{5}$$

$$= \frac{129}{5}$$

$$= 25.8$$

Thus, the mean is 25.8.

20.

**Sol:**

Let us choose  $a = 40$ ,  $h = 10$ , then  $d_i = x_i - 40$  and  $u_i = \frac{x_i - 40}{10}$

Using step-deviation method, the given data is shown as follows:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 40$	$u_i = \frac{x_i - 40}{10}$	$(f_i u_i)$
5 – 15	6	10	-30	-3	-18
15 – 25	10	20	-20	-2	-20
25 – 35	16	30	-10	-1	-16

35 – 45	15	40	0	0	0
45 – 55	24	50	10	1	24
55 – 65	8	60	20	2	16
65 – 75	7	70	30	3	21
Total	$\Sigma f_i = 86$				$\Sigma f_i u_i = 7$

The mean of the data is given by,

$$\begin{aligned}\bar{x} &= a + \left( \frac{\sum_i f_i u_i}{\sum_i f_i} \right) \times h \\ &= 40 + \frac{7}{86} \times 10 \\ &= 40 + \frac{70}{86} \\ &= 40 + 0.81 \\ &= 40.81\end{aligned}$$

21.

**Sol:**

Let us choose  $a = 202.5$ ,  $h = 1$ , then  $d_i = x_i - 202.5$  and  $u_i = \frac{x_i - 202.5}{1}$

Using step-deviation method, the given data is shown as follows:

Weight	Number of packets ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 202.5$	$u_i = \frac{x_i - 202.5}{1}$	$(f_i u_i)$
200 - 201	13	200.5	-2	-2	-26
201 – 202	27	201.5	-1	-1	-27
202 – 203	18	202.5	0	0	0
203 – 204	10	203.5	1	1	10
204 – 205	1	204.5	2	2	2
205 – 206	1	205.5	3	3	3
Total	$\Sigma f_i = 70$				$\Sigma f_i u_i = -38$

The mean of the given data is given by,

$$\begin{aligned}\bar{x} &= a + \left( \frac{\sum_i f_i u_i}{\sum_i f_i} \right) \times h \\ &= 202.5 + \left( \frac{-38}{70} \right) \times 1 \\ &= 202.5 - 0.542 \\ &= 201.96\end{aligned}$$

Hence, the mean is 201.96 g.

22.

**Sol:**

Let us choose  $a = 45$ ,  $h = 10$ , then  $d_i = x_i - 45$  and  $u_i = \frac{x_i - 45}{10}$

Using step-deviation method, the given data is shown as follows:

Weight	Number of packets ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 45$	$u_i = \frac{x_i - 45}{10}$	$(f_i u_i)$
20 – 30	25	35	-20	-2	-50
30 – 40	40	35	-10	-1	-40
40 – 50	42	45	0	0	0
50 – 60	33	55	10	1	33
60 – 70	10	65	20	2	20
Total	$\Sigma f_i = 150$				$\Sigma f_i u_i = -37$

The mean of the given data is given by,

$$\begin{aligned} \bar{x} &= a + \left( \frac{\sum f_i u_i}{\sum f_i} \right) \times h \\ &= 45 - \left( \frac{37}{150} \right) \times 10 \\ &= 45 - \frac{37}{15} \\ &= 45 - 2.466 \\ &= 42.534 \end{aligned}$$

Hence, the mean is 42.534.

23.

Find the mean marks.

**Sol:**

Let us choose  $a = 52.5$ ,  $h = 15$ , then  $d_i = x_i - 52.5$  and  $u_i = \frac{x_i - 52.5}{15}$

Using step-deviation method, the given data is shown as follows:

Weight	Number of students ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 37.5$	$u_i = \frac{x_i - 52.5}{15}$	$(f_i u_i)$
0 – 15	2	7.5	-45	-3	-6
15 – 30	4	22.5	-30	-2	-8
30 – 45	5	37.5	-15	-1	-5

45 – 60	20	52.5	0	0	0
60 – 75	9	67.5	15	1	9
75 – 90	10	82.5	30	2	20
Total	$\Sigma f_i = 50$				$\Sigma f_i u_i = 10$

The mean of the given data is given by,

$$\begin{aligned}\bar{x} &= a + \left( \frac{\sum_i f_i u_i}{\sum_i f_i} \right) \times h \\ &= 52.5 + \left( \frac{10}{50} \right) \times 15 \\ &= 52.5 + 3 \\ &= 55.5\end{aligned}$$

Thus, the mean is 55.5.

24.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$u_i = \frac{(x_i - A)}{h}$ $= \frac{(x_i - 33)}{6}$	$(f_i \times u_i)$
18 – 24	6	21	-2	-12
24 – 30	8	27	-1	-8
30 – 36	12	33 = A	0	0
36 – 42	8	39	1	8
42 – 48	4	45	2	8
48 – 54	2	51	3	6
Total	$\Sigma f_i = 40$			$\Sigma (f_i \times u_i) = 2$

Now,  $A = 33$ ,  $h = 6$ ,  $\Sigma f_i = 40$  and  $\Sigma (f_i \times u_i) = 2$

$$\begin{aligned}\therefore \text{Mean, } \bar{x} &= A + \left\{ h \times \frac{\sum (f_i \times u_i)}{\sum f_i} \right\} \\ &= 33 + \left\{ 6 \times \frac{2}{40} \right\} \\ &= 33 + 0.3 \\ &= 33.3\end{aligned}$$

$\therefore \bar{x} = 33.3$  years

25.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$u_i = \frac{(x_i - A)}{h}$ $= \frac{(x_i - 550)}{20}$	( $f_i \times u_i$ )
500 – 520	14	510	-2	-28
520 – 540	9	530	-1	-9
540 – 560	5	550 = A	0	0
560 – 580	4	570	1	4
580 – 600	3	590	2	6
600 – 620	5	610	3	15
	$\Sigma f_i = 40$			$\Sigma (f_i \times u_i) = -12$

Now,  $A = 550$ ,  $h = 20$ ,  $\Sigma f_i = 40$  and  $\Sigma (f_i \times u_i) = -12$

$$\therefore \text{Mean, } \bar{x} = A + \left\{ h \times \frac{\Sigma (f_i \times u_i)}{\Sigma f_i} \right\}$$

$$= 550 + \left\{ 20 \times \frac{(-12)}{40} \right\}$$

$$= 550 - 6$$

$$= 544$$

$$\therefore \bar{x} = 544$$

26.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$u_i = \frac{(x_i - A)}{h}$ $= \frac{(x_i - 42)}{5}$	( $f_i \times u_i$ )
24.5 – 29.5	4	27	-3	-12
29.5 – 34.5	14	32	-2	-28
34.5 – 39.5	22	37	-1	-22
39.5 – 44.5	16	42 = A	0	0
44.5 – 49.5	6	47	1	6
49.5 – 54.5	5	52	2	10
54.5 – 59.5	3	57	3	9
	$\Sigma f_i = 70$			$\Sigma (f_i \times u_i) = -37$

Now,  $A = 42$ ,  $h = 5$ ,  $\Sigma f_i = 70$  and  $\Sigma (f_i \times u_i) = -37$



$$\begin{aligned} \therefore \text{Mean, } \bar{x} &= A + \left\{ h \times \frac{\sum (f_i \times u_i)}{\sum f_i} \right\} \\ &= 42 + \left\{ 5 \times \frac{(-37)}{70} \right\} \\ &= 42 - 2.64 \\ &= 39.36 \\ \therefore \bar{x} &= 39.36 \\ \therefore \text{Mean age} &= 39.36 \text{ years.} \end{aligned}$$

27.

**Sol:**

Class	Frequency ( $f_i$ )	Mid values ( $x_i$ )	$u_i = \frac{(x_i - A)}{h}$ $= \frac{(x_i - 29.5)}{10}$	( $f_i \times u_i$ )
4.5 – 14.5	6	9.5	-2	-12
14.5 – 24.5	11	19.5	-1	-11
24.5 – 34.5	21	29.5 = A	0	0
34.5 – 44.5	23	39.5	1	23
44.5 – 54.5	14	49.5	2	28
54.5 – 64.5	5	59.5	3	15
	$\Sigma f_i = 80$			$\Sigma (f_i \times u_i) = 43$

Now,  $A = 29.5$ ,  $h = 10$ ,  $\Sigma f_i = 80$  and  $\Sigma (f_i \times u_i) = 43$

$$\begin{aligned} \therefore \text{Mean, } \bar{x} &= A + \left\{ h \times \frac{\sum (f_i \times u_i)}{\sum f_i} \right\} \\ &= 29.5 + \left\{ 10 \times \frac{43}{80} \right\} \\ &= 29.5 + 5.375 \\ &= 34.875 \\ \therefore \bar{x} &= 34.875 \end{aligned}$$

$\therefore$  The average age of the patients is 34.87 years.

28.

**Sol:**

Let us choose  $a = 92$ ,  $h = 5$ , then  $d_i = x_i - 92$  and  $u_i = \frac{x_i - 92}{5}$

Using step-deviation method, the given data is shown as follows:

Weight (in grams)	Number of eggs ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 92$	$u_i = \frac{x_i - 92}{5}$	$(f_i u_i)$
74.5 – 79.5	4	77	-15	-3	-12
79.5 – 84.5	9	82	-10	-2	-18
84.5 – 89.5	13	87	-5	-1	-13
89.5 – 94.5	17	92	0	0	0
94.5 – 99.5	12	97	5	1	12
99.5 – 104.5	3	102	10	2	6
104.5 – 109.5	2	107	15	3	6
Total	$\Sigma f_i = 60$				$\Sigma f_i u_i = -19$

The mean of the given data is given by,

$$\begin{aligned} \bar{x} &= a + \left( \frac{\sum f_i u_i}{\sum f_i} \right) \times h \\ &= 92 + \left( \frac{-19}{60} \right) \times 5 \\ &= 92 - 1.58 \\ &= 90.42 \\ &\approx 90 \end{aligned}$$

Thus, the mean weight to the nearest gram is 90 g.

29.

**Sol:**

Let us choose  $a = 17.5$ ,  $h = 5$ , then  $d_i = x_i - 17.5$  and  $u_i = \frac{x_i - 17.5}{5}$

Using step-deviation method, the given data is shown as follows:

Marks	Number of students (cf)	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$d_i = x_i - 17.5$	$u_i = \frac{x_i - 17.5}{5}$	$(f_i u_i)$
0 – 5	3	3	2.5	-15	-3	-9
5 – 10	10	7	7.5	-10	-2	-14
10 – 15	25	15	12.5	-5	-1	-15
15 – 20	49	24	17.5	0	0	0
20 – 25	65	16	22.5	5	1	16
25 – 30	73	8	27.5	10	2	16
30 – 35	78	5	32.5	15	3	15
35 – 40	80	2	37.5	20	4	8

Total		$\Sigma f_i = 80$				$\Sigma f_i u_i = 17$
-------	--	-------------------	--	--	--	-----------------------

The mean of the given data is given by,

$$\begin{aligned}\bar{x} &= a + \left( \frac{\sum_i f_i u_i}{\sum_i f_i} \right) \times h \\ &= 17.5 + \left( \frac{17}{80} \right) \times 5 \\ &= 17.5 + 1.06 \\ &= 18.56\end{aligned}$$

Thus, the mean marks correct to 2 decimal places is 18.56.

### Exercise 9B

1.

**Sol:**

We prepare the cumulative frequency table, as shown below:

Age (in years)	Number of patients ( $f_i$ )	Cumulative Frequency (cf)
0 – 15	5	5
15 – 30	20	25
30 – 45	40	65
45 – 60	50	115
60 – 75	25	140
Total	$N = \Sigma f_i = 140$	

Now,  $N = 140 \Rightarrow \frac{N}{2} = 70$ .

The cumulative frequency just greater than 70 is 115 and the corresponding class is 45 – 60.

Thus, the median class is 45 – 60.

$\therefore l = 45, h = 15, f = 50, N = 140$  and  $cf = 65$ .

Now,

$$\begin{aligned}\text{Median} &= l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h \\ &= 45 + \left( \frac{\frac{140}{2} - 65}{50} \right) \times 15 \\ &= 45 + \left( \frac{70 - 65}{50} \right) \times 15 \\ &= 45 + 1.5 \\ &= 46.5\end{aligned}$$

Hence, the median age is 46.5 years.

2.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
0 – 7	3	3
7 – 14	4	7
14 – 21	7	14
21 – 28	11	25
28 – 35	0	25
35 – 42	16	41
42 – 49	9	50
	$N = \sum f = 50$	

Now,  $N = 50 \Rightarrow \frac{N}{2} = 25$ .

The cumulative frequency just greater than 25 is 41 and the corresponding class is 35 – 42. Thus, the median class is 35 – 42.

$\therefore l = 35, h = 7, f = 16, cf = \text{c.f. of preceding class} = 25$  and  $\frac{N}{2} = 25$ .

Now,

$$\begin{aligned} \text{Median} &= l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h \\ &= 35 + 7 \times \left( \frac{25 - 25}{16} \right) \\ &= 35 + 0 \\ &= 35 \end{aligned}$$

Hence, the median age is 46.5 years.

3.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
0 – 100	40	40
100 – 200	32	72
200 – 300	48	120
300 – 400	22	142
400 – 500	8	150
	$N = \sum f = 150$	

Now,  $N = 150$

$$\Rightarrow \frac{N}{2} = 75.$$

The cumulative frequency just greater than 75 is 120 and the corresponding class is 200 – 300.

Thus, the median class is 200 – 300.

$$\therefore l = 200, h = 100, f = 48, cf = \text{c.f. of preceding class} = 72 \text{ and } \frac{N}{2} = 75.$$

Now,

$$\begin{aligned} \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 200 + \left\{ 100 \times \left( \frac{75 - 72}{48} \right) \right\} \\ &= 200 + 6.25 \\ &= 206.25 \end{aligned}$$

Hence, the median daily wage income of the workers is Rs 206.25.

4.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
5 – 10	5	5
10 – 15	6	11
15 – 20	15	26
20 – 25	10	36
25 – 30	5	41
30 – 35	4	45
35 – 40	2	47
40 – 45	2	49
	$N = \sum f = 49$	

Now,  $N = 49$

$$\Rightarrow \frac{N}{2} = 24.5.$$

The cumulative frequency just greater than 24.5 is 26 and the corresponding class is 15 – 20.

Thus, the median class is 15 – 20.

$$\therefore l = 15, h = 5, f = 15, cf = \text{c.f. of preceding class} = 11 \text{ and } \frac{N}{2} = 24.5.$$

Now,

$$\begin{aligned} \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 15 + \left\{ 5 \times \left( \frac{24.5 - 11}{15} \right) \right\} \end{aligned}$$

$$= 15 + 4.5$$

$$= 19.5$$

Hence, the median = 19.5.

5.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
65- 85	4	4
85 – 105	5	9
105 – 125	13	22
125 – 145	20	42
145 – 165	14	56
165 – 185	7	63
185 – 205	4	67
	$N = \sum f = 67$	

Now,  $N = 67$

$$\Rightarrow \frac{N}{2} = 33.5.$$

The cumulative frequency just greater than 33.5 is 42 and the corresponding class is 125 - 145.

Thus, the median class is 125 – 145.

$\therefore l = 125, h = 20, f = 20, cf = \text{c.f. of preceding class} = 22$  and  $\frac{N}{2} = 33.5$ .

Now,

$$\begin{aligned} \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 125 + \left\{ 20 \times \left( \frac{33.5 - 22}{20} \right) \right\} \\ &= 125 + 11.5 \\ &= 136.5 \end{aligned}$$

Hence, the median = 136.5.

6.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
135 – 140	6	6
140 – 145	10	16
145 – 150	18	34
150 – 155	22	56
155 – 160	20	76
160 – 165	15	91
165 – 170	6	97
170 – 175	3	100
	$N = \sum f = 100$	

Now,  $N = 100$

$$\Rightarrow \frac{N}{2} = 50.$$

The cumulative frequency just greater than 50 is 56 and the corresponding class is 150 - 155.

Thus, the median class is 150 – 155.

$$\therefore l = 150, h = 5, f = 22, cf = \text{c.f. of preceding class} = 34 \text{ and } \frac{N}{2} = 50.$$

Now,

$$\begin{aligned} \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 150 + \left\{ 5 \times \left( \frac{50 - 34}{22} \right) \right\} \\ &= 150 + 3.64 \\ &= 153.64 \end{aligned}$$

Hence, the median = 153.64.

7.

**Sol:**

Class	Frequency (f <sub>i</sub> )	Cumulative Frequency (cf)
0 – 10	5	5
10 – 20	25	30
20 – 30	x	x + 30
30 – 40	18	x + 48
40 – 50	7	x + 55

Median is 24 which lies in 20 – 30

$\therefore$  Median class = 20 – 30

Let the unknown frequency be x.

Here,  $l = 20$ ,  $\frac{n}{2} = \frac{x+55}{2}$ , c.f. of the preceding class = c.f = 30,  $f = x$ ,  $h = 10$

Now,

$$\text{Median, } M = l + \frac{\frac{n}{2} - cf}{f} \times h$$

$$\Rightarrow 24 = 20 + \frac{\frac{x+55}{2} - 30}{x} \times 10$$

$$\Rightarrow 24 = 20 + \frac{x+55-60}{x} \times 10$$

$$\Rightarrow 24 = 20 + \frac{x-5}{x} \times 10$$

$$\Rightarrow 24 = 20 + \frac{5x-25}{x}$$

$$\Rightarrow 24 = \frac{20+5x-25}{x}$$

$$\Rightarrow 24x = 25x - 25$$

$$\Rightarrow -x = -25$$

$$\Rightarrow x = 25$$

Hence, the unknown frequency is 25.

8.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
0 – 5	12	12
5 – 10	a	12 + a
10 – 15	12	24 + a
15 – 20	15	39 + a
20 – 25	b	39 + a + b
25 – 30	6	45 + a + b
30 – 35	6	51 + a + b
35 – 40	4	55 + a + b
Total	$N = \sum f_i = 70$	

Let a and b be the missing frequencies of class intervals 5 – 10 and 20 – 25 respectively.

$$\text{Then, } 55 + a + b = 70 \Rightarrow a + b = 15 \dots (1)$$

Median is 16, which lies in 15 – 20. So, the median class is 15 – 20.

$$\therefore l = 15, h = 5, N = 70, f = 15 \text{ and } cf = 24 + a$$

Now,

$$\text{Median, } M = l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h$$



$$\Rightarrow 16 = 15 + \left( \frac{\frac{70}{2} - (24 + a)}{15} \right) \times 5$$

$$\Rightarrow 16 = 15 + \left( \frac{35 - 24 - a}{3} \right)$$

$$\Rightarrow 16 = 15 + \left( \frac{11 - a}{3} \right)$$

$$\Rightarrow 16 - 15 = \frac{11 - a}{3}$$

$$\Rightarrow 1 \times 3 = 11 - a$$

$$\Rightarrow a = 11 - 3$$

$$\Rightarrow a = 8$$

$$\therefore b = 15 - a \quad [\text{From (1)}]$$

$$\Rightarrow b = 15 - 8$$

$$\Rightarrow b = 7$$

Hence,  $a = 8$  and  $b = 7$ .

9.

**Sol:**

We prepare the cumulative frequency table, as shown below:

Runs scored	Number of batsman ( $f_i$ )	Cumulative Frequency (cf)
2500 – 3500	5	5
3500 – 4500	x	5 + x
4500 – 5500	y	5 + x + y
5500 – 6500	12	17 + x + y
6500 – 7500	6	23 + x + y
7500 – 8500	2	25 + x + y
Total	$N = \sum f_i = 60$	

Let x and y be the missing frequencies of class intervals 3500 – 4500 respectively. Then,

$$25 + x + y = 60 \Rightarrow x + y = 35 \quad \dots\dots(1)$$

Median is 5000, which lies in 4500 – 5500. So, the median class is 4500 – 5500.

$$\therefore l = 4500, h = 1000, N = 60, f = y \text{ and } cf = 5 + x$$

Now,

$$\text{Median, } M = l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h$$

$$\Rightarrow 5000 = 4500 + \left( \frac{\frac{60}{2} - (5+x)}{y} \right) \times 1000$$

$$\Rightarrow 5000 - 4500 = \left( \frac{30-5-x}{y} \right) \times 1000$$

$$\Rightarrow 500 = \left( \frac{25-x}{y} \right) \times 1000$$

$$\Rightarrow y = 50 - 2x$$

$$\Rightarrow 35 - x = 50 - 2x \quad [\text{From (1)}]$$

$$\Rightarrow 2x - x = 50 - 35$$

$$\Rightarrow x = 15$$

$$\therefore y = 35 - x \quad [\text{From (1)}]$$

$$\Rightarrow y = 35 - 15$$

$$\Rightarrow y = 20$$

Hence,  $x = 15$  and  $y = 20$ .

10.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
0 - 10	$f_1$	$f_1$
10 - 20	5	$f_1 + 5$
20 - 30	9	$f_1 + 14$
30 - 40	12	$f_1 + 26$
40 - 50	$f_2$	$f_1 + f_2 + 26$
50 - 60	3	$f_1 + f_2 + 29$
60 - 70	2	$f_1 + f_2 + 31$
	$N = \sum f = 40$	

Now,  $f_1 + f_2 + 31 = 40$

$$\Rightarrow f_1 + f_2 = 9$$

$$\Rightarrow f_2 = 9 - f_1$$

The median is 32.5 which lies in 30 - 40.

Hence, median class = 30 - 40

Here,  $l = 30$ ,  $\frac{N}{2} = \frac{40}{2} = 20$ ,  $f = 12$  and  $cf = 14 + f_1$

Now, median = 32.5

$$\Rightarrow l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h = 32.5$$

$$\Rightarrow 30 + \left( \frac{20 - (14 + f_1)}{12} \right) \times 10 = 32.5$$

$$\Rightarrow \frac{6 - f_1}{12} \times 10 = 2.5$$

$$\Rightarrow \frac{60 - 10f_1}{12} = 2.5$$

$$\Rightarrow 60 - 10f_1 = 30$$

$$\Rightarrow 10f_1 = 30$$

$$\Rightarrow f_1 = 3$$

From equation (i), we have:

$$f_2 = 9 - 3$$

$$\Rightarrow f_2 = 6$$

11.

**Sol:** First, we will convert the data into exclusive form.

Class	Frequency (f)	Cumulative Frequency (cf)
18.5 – 25.5	35	35
25.5 – 32.5	96	131
32.5 – 39.5	68	199
39.5 – 46.5	102	301
46.5 – 53.5	35	336
53.5 – 60.5	4	340
	$N = \sum f = 340$	

Now,  $N = 340$

$$\Rightarrow \frac{N}{2} = 170.$$

The cumulative frequency just greater than 170 is 199 and the corresponding class is 32.5 – 39.5.

Thus, the median class is 32.5 – 39.5.

$\therefore l = 32.5$ ,  $h = 7$ ,  $f = 68$ ,  $cf = \text{c.f. of preceding class} = 131$  and  $\frac{N}{2} = 170$ .

$$\begin{aligned} \therefore \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 32.5 + \left\{ 7 \times \left( \frac{170 - 131}{68} \right) \right\} \\ &= 32.5 + 4.01 \\ &= 36.51 \end{aligned}$$

Hence, the median = 36.51.

12.

**Sol:**

Class	Frequency (f)	Cumulative Frequency (cf)
60.5 – 70.5	5	5
70.5 – 80.5	15	20
80.5 – 90.5	20	40
90.5 – 100.5	30	70
100.5 – 110.5	20	90
110.5 – 120.5	8	98
	$N = \sum f = 98$	

Now,  $N = 98$ 

$$\Rightarrow \frac{N}{2} = 49.$$

The cumulative frequency just greater than 49 is 70 and the corresponding class is 90.5 – 100.5.

Thus, the median class is 90.5 – 100.5.

Now,  $l = 90.5$ ,  $h = 10$ ,  $f = 30$ ,  $cf = \text{c.f. of preceding class} = 40$  and  $\frac{N}{2} = 49$ .

$$\begin{aligned} \therefore \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 90.5 + \left\{ 10 \times \left( \frac{49 - 40}{30} \right) \right\} \\ &= 90.5 + 3 \\ &= 93.5 \end{aligned}$$

Hence, median wages = Rs. 93.50.

13.

**Sol:**

Converting into exclusive form, we get:

Class	Frequency (f)	Cumulative Frequency (cf)
0.5 – 5.5	7	7
5.5 – 10.5	10	17
10.5 – 15.5	16	33
15.5 – 20.5	32	65
20.5 – 25.5	24	89
25.5 – 30.5	16	105
30.5 – 35.5	11	116

35.5 – 40.5	5	121
40.5 – 45.5	2	123
	$N = \sum f = 123$	

Now,  $N = 123$

$$\Rightarrow \frac{N}{2} = 61.5.$$

The cumulative frequency just greater than 61.5 is 65 and the corresponding class is 15.5 – 20.5.

Thus, the median class is 15.5 – 20.5.

$\therefore l = 15.5$ ,  $h = 5$ ,  $f = 32$ ,  $cf = \text{c.f. of preceding class} = 33$  and  $\frac{N}{2} = 61.5$ .

$$\begin{aligned} \therefore \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 15.5 + \left\{ 5 \times \left( \frac{61.5 - 33}{32} \right) \right\} \\ &= 15.5 + 4.45 \\ &= 19.95 \end{aligned}$$

Hence, median = 19.95.

14.

**Sol:**

Class	Cumulative frequency (cf)	Frequency (f)
0 – 10	12	12
10 – 20	32	20
20 – 30	57	25
30 – 40	80	23
40 – 50	92	12
50 – 60	116	24
60 – 70	164	48
70 – 80	200	36
		$N = \sum f = 200$

Now,  $N = 200$

$$\Rightarrow \frac{N}{2} = 100.$$

The cumulative frequency just greater than 100 is 116 and the corresponding class is 50 – 60.

Thus, the median class is 50 – 60.

$\therefore l = 50, h = 10, f = 24, cf = \text{c.f. of preceding class} = 92$  and  $\frac{N}{2} = 100$ .

$$\begin{aligned}\therefore \text{Median, } M &= l + \left\{ h \times \left( \frac{\frac{N}{2} - cf}{f} \right) \right\} \\ &= 50 + \left\{ 10 \times \left( \frac{100 - 92}{24} \right) \right\} \\ &= 50 + 3.33 \\ &= 53.33\end{aligned}$$

Hence, median = 53.33.

### Exercise 9C

1.

**Sol:**

Here, the maximum class frequency is 45, and the class corresponding to this frequency is 30 – 40. So, the modal class is 30- 40.

Now,

Modal class = 30 – 40, lower limit ( $l$ ) of modal class = 30, class size ( $h$ ) = 10,

frequency ( $f_1$ ) of the modal class = 45,

frequency ( $f_0$ ) of class preceding the modal class = 35,

frequency ( $f_2$ ) of class succeeding the modal class = 25

Now, let us substitute these values in the formula:

$$\begin{aligned}\text{Mode} &= l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\ &= 30 + \left( \frac{45 - 35}{90 - 35 - 45} \right) \times 10 \\ &= 30 + \left( \frac{10}{30} \right) \times 10 \\ &= 30 + 3.33 \\ &= 33.33\end{aligned}$$

Hence, the mode is 33.33.

2.

**Sol:**

Here, the maximum class frequency is 28, and the class corresponding to this frequency is 40 – 60. So, the modal class is 40 – 60.

Now,

Modal class = 40 – 60, lower limit ( $l$ ) of modal class = 40, class size ( $h$ ) = 20,

frequency ( $f_1$ ) of the modal class = 28,

frequency ( $f_0$ ) of class preceding the modal class = 16,

frequency ( $f_2$ ) of class succeeding the modal class = 20

Now, let us substitute these values in the formula:

$$\begin{aligned}\text{Mode} &= l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\ &= 40 + \left( \frac{28 - 16}{56 - 16 - 20} \right) \times 20 \\ &= 40 + \left( \frac{12}{20} \right) \times 20 \\ &= 40 + 12 \\ &= 52\end{aligned}$$

Hence, the mode is 52.

3.

**Sol:**

Here, the maximum class frequency is 20, and the class corresponding to this frequency is 160 – 165. So, the modal class is 160 – 165.

Now,

Modal class = 160 – 165, lower limit ( $l$ ) of modal class = 160, class size ( $h$ ) = 5,

frequency ( $f_1$ ) of the modal class = 20,

frequency ( $f_0$ ) of class preceding the modal class = 8,

frequency ( $f_2$ ) of class succeeding the modal class = 12

Now, let us substitute these values in the formula:

$$\begin{aligned}\text{Mode} &= l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\ &= 160 + \left( \frac{20 - 8}{40 - 8 - 12} \right) \times 5 \\ &= 160 + \left( \frac{12}{20} \right) \times 5 \\ &= 160 + 3 \\ &= 163\end{aligned}$$

Hence, the mode is 163.

It represents that the height of maximum number of students is 163cm.

Now, to find the mean let us put the data in the table given below:

Height (in cm)	Number of students ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
150 – 155	15	152.5	2287.5
155 – 160	8	157.5	1260
160 – 165	20	162.5	3250
165 – 170	12	167.5	2010
170 – 175	5	172.5	862.5
Total	$\Sigma f_i = 60$		$\Sigma f_i x_i = 9670$

$$\begin{aligned} \text{Mean} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ &= \frac{9670}{60} \\ &= 161.17 \end{aligned}$$

Thus, mean of the given data is 161.17.

It represents that on an average, the height of a student is 161.17cm.

4.

**Sol:**

As the class 26 – 30 has the maximum frequency, it is the modal class.

Now,  $x_k = 26$ ,  $h = 4$ ,  $f_k = 25$ ,  $f_{k-1} = 20$ ,  $f_{k+1} = 22$

$$\begin{aligned} \therefore \text{Mode, } M_0 &= x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\} \\ &= 26 + \left\{ 4 \times \frac{(25-20)}{(2 \times 25 - 20 - 22)} \right\} \\ &= 26 + \left\{ 4 \times \frac{5}{8} \right\} \\ &= (26 + 2.5) \\ &= 28.5 \end{aligned}$$



5.

**Sol:**

As the class 1500-2000 has the maximum frequency, it is the modal class.

Now,  $x_k = 1500$ ,  $h = 500$ ,  $f_k = 40$ ,  $f_{k-1} = 24$ ,  $f_{k+1} = 31$

$$\therefore \text{Mode, } M_0 = x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\}$$

$$= 1500 + \left\{ 500 \times \frac{(40 - 24)}{(2 \times 40 - 24 - 31)} \right\}$$

$$= 1500 + \left\{ 500 \times \frac{16}{25} \right\}$$

$$= (1500 + 320)$$

$$= 1820$$

Hence, mode = Rs 1820

6.

**Sol:**

As the class 5000-10000 has the maximum frequency, it is the modal class.

Now,  $x_k = 5000$ ,  $h = 5000$ ,  $f_k = 150$ ,  $f_{k-1} = 90$ ,  $f_{k+1} = 100$

$$\therefore \text{Mode, } M_0 = x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\}$$

$$= 5000 + \left\{ 5000 \times \frac{(150 - 90)}{(2 \times 150 - 90 - 100)} \right\}$$

$$= 5000 + \left\{ 5000 \times \frac{60}{110} \right\}$$

$$= (5000 + 2727.27)$$

$$= 7727.27$$

Hence, mode = Rs 7727.27

7.

**Sol:**

As the class 15 – 20 has the maximum frequency, it is the modal class.

Now,  $x_k = 15$ ,  $h = 5$ ,  $f_k = 24$ ,  $f_{k-1} = 18$ ,  $f_{k+1} = 17$

$$\begin{aligned}
 \therefore \text{Mode, } M_0 &= x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\} \\
 &= 15 + \left\{ 5 \times \frac{(24-18)}{(2 \times 24 - 18 - 17)} \right\} \\
 &= 15 + \left\{ 5 \times \frac{6}{13} \right\} \\
 &= (15 + 2.3) \\
 &= 17.3 \\
 \text{Hence, mode} &= 17.3 \text{ years}
 \end{aligned}$$

8.

**Sol:**

As the class 85 – 95 has the maximum frequency, it is the modal class.

Now,  $x_k = 85$ ,  $h = 10$ ,  $f_k = 32$ ,  $f_{k-1} = 30$ ,  $f_{k+1} = 6$

$$\begin{aligned}
 \therefore \text{Mode, } M_0 &= x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\} \\
 &= 85 + \left\{ 10 \times \frac{(32-30)}{(2 \times 32 - 30 - 6)} \right\} \\
 &= 85 + \left\{ 10 \times \frac{2}{28} \right\} \\
 &= (85 + 0.71) \\
 &= 85.71 \\
 \text{Hence, mode} &= 85.71
 \end{aligned}$$

9.

**Sol:**

Clearly, we have to find the mode of the data. The given data is an inclusive series. So, we will convert it to an exclusive form as given below:

Class interval	0.5 – 5.5	5.5 – 10.5	10.5 – 15.5	15.5 – 20.5	20.5 – 25.5	25.5 – 30.5	30.5 – 35.5	35.5 – 40.5	40.5 – 45.5	45.5 – 50.5
Frequency	3	8	13	18	28	20	13	8	6	4

As the class 20.5 – 25.5 has the maximum frequency, it is the modal class.

Now,  $x_k = 20.5$ ,  $h = 5$ ,  $f_k = 28$ ,  $f_{k-1} = 18$ ,  $f_{k+1} = 20$

$$\begin{aligned}
 \therefore \text{Mode, } M_0 &= x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\} \\
 &= 20.5 + \left\{ 5 \times \frac{(28-18)}{(2 \times 28 - 18 - 20)} \right\} \\
 &= 20.5 + \left\{ 5 \times \frac{10}{18} \right\} \\
 &= (20.5 + 2.78)
 \end{aligned}$$

$$= 23.28$$

Hence, mode = 23.28

10.

**Sol:**

It is given that the sum of frequencies is 181.

$$\therefore x + 15 + 18 + 30 + 50 + 48 + x = 181$$

$$\Rightarrow 2x + 161 = 181$$

$$\Rightarrow 2x = 181 - 161$$

$$\Rightarrow 2x = 20$$

$$\Rightarrow x = 10$$

Thus,  $x = 10$

Here, the maximum class frequency is 50, and the class corresponding to this frequency is 13 – 15. So, the modal class is 13 – 15.

Now,

Modal class = 13 – 15, lower limit ( $l$ ) of modal class = 13, class size ( $h$ ) = 2,

frequency ( $f_1$ ) of the modal class = 50,

frequency ( $f_0$ ) of class preceding the modal class = 30,

frequency ( $f_2$ ) of class succeeding the modal class = 48

Now, let us substitute these values in the formula:

$$\begin{aligned} \text{Mode} &= l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h \\ &= 13 + \left( \frac{50 - 30}{100 - 30 - 48} \right) \times 2 \\ &= 13 + \left( \frac{20}{22} \right) \times 2 \\ &= 13 + 1.82 \\ &= 14.82 \end{aligned}$$

Hence, the mode is 14.82.

## Exercise 9D

1.

**Sol:**

To find the mean let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
0 – 10	4	5	20
10 – 20	4	15	60
20 – 30	7	25	175
30 – 40	10	35	350
40 – 50	12	45	540
50 – 60	8	55	440
60 – 70	5	65	325
Total	$\sum f_i = 50$		$\sum f_i x_i = 1910$

$$\begin{aligned} \text{Mean} &= \frac{\sum f_i x_i}{\sum f_i} \\ &= \frac{1910}{50} \\ &= 38.2 \end{aligned}$$

Thus, the mean of the given data is 38.2.

Now, to find the median let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative Frequency (cf)
0 – 10	4	4
10 – 20	4	8
20 – 30	7	15
30 – 40	10	25
40 – 50	12	37
50 – 60	8	45
60 – 70	5	50
Total	$N = \sum f_i = 50$	

$$\text{Now, } N = 50 \Rightarrow \frac{N}{2} = 25.$$

The cumulative frequency just greater than 25 is 37 and the corresponding class is 40 – 50.

Thus, the median class is 40 – 50.

$$\therefore l = 40, h = 10, N = 50, f = 12 \text{ and } cf = 25.$$

Now,

$$\text{Median} = l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h$$

$$= 40 + \left(\frac{25 - 25}{12}\right) \times 10$$

$$= 40$$

Thus, the median is 40.

We know that,

$$\text{Mode} = 3(\text{median}) - 2(\text{mean})$$

$$= 3 \times 40 - 2 \times 38.2$$

$$= 120 - 76.4$$

$$= 43.6$$

Hence, Mean = 38.2, Median = 40 and Mode = 43.6

2.

**Sol:**

To find the mean let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
0 – 20	6	10	60
20 – 40	8	30	240
40 – 60	10	50	500
60 – 80	12	70	840
80 – 100	6	90	540
100 – 120	5	110	550
120 – 140	3	130	390
Total	$\Sigma f_i = 50$		$\Sigma f_i x_i = 3120$

$$\text{Mean} = \frac{\sum f_i x_i}{\sum f_i}$$

$$= \frac{3120}{50}$$

$$= 62.4$$

Thus, the mean of the given data is 62.4.

Now, to find the median let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative Frequency (cf)
0 – 20	6	6
20 – 40	8	14
40 – 60	10	24
60 – 80	12	36
80 – 100	6	42
100 – 120	5	47
120 – 140	3	50
Total	$N = \Sigma f_i = 50$	

$$\text{Now, } N = 50 \Rightarrow \frac{N}{2} = 25.$$

The cumulative frequency just greater than 25 is 36 and the corresponding class is 60 – 80.  
Thus, the median class is 60 – 80.

$$\therefore l = 60, h = 20, N = 50, f = 12 \text{ and } cf = 24.$$

Now,

$$\begin{aligned} \text{Median} &= l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h \\ &= 60 + \left( \frac{25 - 24}{12} \right) \times 20 \\ &= 60 + 1.67 \\ &= 61.67 \end{aligned}$$

Thus, the median is 61.67.

We know that,

$$\begin{aligned} \text{Mode} &= 3(\text{median}) - 2(\text{mean}) \\ &= 3 \times 61.67 - 2 \times 62.4 \\ &= 185.01 - 124.8 \\ &= 60.21 \end{aligned}$$

Hence, Mean = 62.4, Median = 61.67 and Mode = 60.21

3.

**Sol:**

To find the mean let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
0 – 50	2	25	50
50 – 100	3	75	225
100 – 150	5	125	625
150 – 200	6	175	1050
200 – 250	5	225	1125
250 – 300	3	275	825
300 – 350	1	325	325
Total	$\sum f_i = 25$		$\sum f_i x_i = 4225$

$$\begin{aligned} \text{Mean} &= \frac{\sum f_i x_i}{\sum f_i} \\ &= \frac{4225}{25} \\ &= 169 \end{aligned}$$

Thus, mean of the given data is 169.

Now, to find the median let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative Frequency (cf)
0 – 50	2	2
50 – 100	3	5
100 – 150	5	10
150 – 200	6	16
200 – 250	5	21
250 – 300	3	24
300 – 350	1	25
Total	$N = \sum f_i = 25$	

Now,  $N = 25 \Rightarrow \frac{N}{2} = 12.5$ .

The cumulative frequency just greater than 12.5 is 16 and the corresponding class is 150 – 200.

Thus, the median class is 150 – 200.

$\therefore l = 150, h = 50, N = 25, f = 6$  and  $cf = 10$ .

Now,

$$\begin{aligned} \text{Median} &= l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h \\ &= 150 + \left( \frac{12.5 - 10}{6} \right) \times 50 \\ &= 150 + 20.83 \\ &= 170.83 \end{aligned}$$

Thus, the median is 170.83.

We know that,

$$\begin{aligned} \text{Mode} &= 3(\text{median}) - 2(\text{mean}) \\ &= 3 \times 170.83 - 2 \times 169 \\ &= 512.49 - 338 \\ &= 174.49 \end{aligned}$$

Hence, Mean = 169, Median = 170.83 and Mode = 174.49

4.

**Sol:**

To find the mean let us put the data in the table given below:

Marks obtained	Number of students ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
25 – 35	7	30	210
35 – 45	31	40	1240

45 – 55	33	50	1650
55 – 65	17	60	1020
65 – 75	11	70	770
75 – 85	1	80	80
Total	$\Sigma f_i = 100$		$\Sigma f_i x_i = 4970$

$$\begin{aligned}\text{Mean} &= \frac{\sum_i f_i x_i}{\sum_i f_i} \\ &= \frac{4970}{100} \\ &= 49.7\end{aligned}$$

Thus, mean of the given data is 49.7.

Now, to find the median let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative Frequency (cf)
25 – 35	7	7
35 – 45	31	38
45 – 55	33	71
55 – 65	17	88
65 – 75	11	99
75 – 85	1	100
Total	$N = \Sigma f_i = 100$	

$$\text{Now, } N = 100 \Rightarrow \frac{N}{2} = 50.$$

The cumulative frequency just greater than 50 is 71 and the corresponding class is 45 – 55.

Thus, the median class is 45 – 55.

$$\therefore l = 45, h = 10, N = 100, f = 33 \text{ and } cf = 38.$$

Now,

$$\begin{aligned}\text{Median} &= l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h \\ &= 45 + \left( \frac{50 - 38}{33} \right) \times 10 \\ &= 45 + 3.64 \\ &= 48.64\end{aligned}$$

Thus, the median is 48.64.

We know that,

$$\begin{aligned}\text{Mode} &= 3(\text{median}) - 2(\text{mean}) \\ &= 3 \times 48.64 - 2 \times 49.70 \\ &= 145.92 - 99.4 \\ &= 46.52\end{aligned}$$

Hence, Mean = 49.70, Median = 48.64 and Mode = 46.52



5.

**Sol:** We have the following

Height in cm	Mid value ( $x_i$ )	Frequency ( $f_i$ )	Cumulative frequency	( $f_i \times x_i$ )
120 – 130	125	2	2	250
130 – 140	135	8	10	1080
140 – 150	145	12	22	1740
150 – 160	155	20	42	3100
160 – 170	165	8	50	1320
		$\Sigma f_i = 50$		$\Sigma f_i \times x_i = 7490$

$$\text{Mean, } \bar{x} = \frac{\sum (f_i \times x_i)}{\sum f_i}$$

$$= \frac{7490}{50}$$

$$= 149.8$$

Now,  $N = 50$

$$\Rightarrow \frac{N}{2} = 25.$$

The cumulative frequency just greater than 25 is 42 and the corresponding class is 150 – 160.

Thus, the median class is 150 – 160.

$\therefore l = 150, h = 10, f = 20, c = \text{cf of preceding class} = 22$  and  $\frac{N}{2} = 25$

Now,

$$\text{Median, } M_e = l + \left\{ h \times \left( \frac{\frac{N}{2} - c}{f} \right) \right\}$$

$$= 150 + \left\{ 10 \times \left( \frac{25 - 22}{20} \right) \right\}$$

$$= \left( 150 + 10 \times \frac{3}{20} \right)$$

$$= 151.5$$

Mode = 3(median) – 2(mean)

$$= 3 \times 151.5 - 2 \times 149.8$$

$$= 154.9$$

6.

**Sol:**

We have the following:

Daily income	Mid value ( $x_i$ )	Frequency ( $f_i$ )	Cumulative frequency	$(f_i \times x_i)$
100 – 120	110	12	12	1320
120 – 140	130	14	26	1820
140 – 160	150	8	34	1200
160 – 180	170	6	40	1020
180 – 200	190	10	50	1900
		$\Sigma f_i = 50$		$\Sigma f_i \times x_i = 7260$

$$\begin{aligned}\text{Mean, } \bar{x} &= \frac{\Sigma f_i \times x_i}{\Sigma f_i} \\ &= \frac{7260}{50} \\ &= 145.2\end{aligned}$$

Now,  $N = 50$ 

$$\Rightarrow \frac{N}{2} = 25.$$

The cumulative frequency just greater than 25 is 26 and the corresponding class is 120 – 140.

Thus, the median class is 120 – 140.

$$\therefore l = 120, h = 20, f = 14, c = \text{cf of preceding class} = 12 \text{ and } \frac{N}{2} = 25$$

Now,

$$\begin{aligned}\text{Median, } M_e &= l + \left\{ h \times \left( \frac{\frac{N}{2} - c}{f} \right) \right\} \\ &= 120 + \left\{ 20 \times \left( \frac{25 - 12}{14} \right) \right\} \\ &= \left( 120 + 20 \times \frac{13}{14} \right) \\ &= 138.57\end{aligned}$$

$$\text{Mode} = 3(\text{median}) - 2(\text{mean})$$

$$\begin{aligned}&= 3 \times 138.57 - 2 \times 145.2 \\ &= 125.31\end{aligned}$$

7.

**Sol:**

We have the following:

Daily expenditure (in Rs)	Mid value ( $x_i$ )	Frequency ( $f_i$ )	Cumulative frequency	( $f_i \times x_i$ )
100 – 150	125	6	6	750
150 – 200	175	7	13	1225
200 – 250	225	12	25	2700
250 – 300	275	3	28	825
300 – 350	325	2	30	650
		$\Sigma f_i = 30$		$\Sigma f_i \times x_i = 6150$

$$\begin{aligned}\text{Mean, } \bar{x} &= \frac{\Sigma f_i \times x_i}{\Sigma f_i} \\ &= \frac{6150}{30} \\ &= 205\end{aligned}$$

Now,  $N = 30$ 

$$\Rightarrow \frac{N}{2} = 15.$$

The cumulative frequency just greater than 15 is 25 and the corresponding class is 200 – 250.

Thus, the median class is 200 – 250.

$$\therefore l = 200, h = 50, f = 12, c = \text{cf of preceding class} = 13 \text{ and } \frac{N}{2} = 15$$

Now,

$$\begin{aligned}\text{Median, } M_e &= l + \left\{ h \times \left( \frac{\frac{N}{2} - c}{f} \right) \right\} \\ &= 200 + \left\{ 50 \times \left( \frac{15 - 13}{12} \right) \right\} \\ &= \left( 200 + 50 \times \frac{2}{12} \right) \\ &= 200 + 8.33 \\ &= 208.33\end{aligned}$$


## Exercise 9E

30.

**Sol:**

The frequency distribution table of less than type is given as follows:


Marks (upper class limits)	Cumulative frequency (cf)
Less than 10	5
Less than 20	$5 + 3 = 8$
Less than 30	$8 + 4 = 12$
Less than 40	$12 + 3 = 15$
Less than 50	$15 + 3 = 18$
Less than 60	$18 + 4 = 22$
Less than 70	$22 + 7 = 29$
Less than 80	$29 + 9 = 38$
Less than 90	$38 + 7 = 45$
Less than 100	$45 + 8 = 53$



Taking upper class limits of class intervals on x-axis and their respective frequencies on y-axis, its ogive can be drawn as follows:

$$\text{Here, } N = 53 \Rightarrow \frac{N}{2} = 26.5.$$

Mark the point A whose ordinate is 26.5 and its x-coordinate is 66.4.




Thus, median of the data is 66.4.

31.


**Sol:**

Taking upper class limits of class intervals on x-axis and their respective frequencies on y-axis, its ogive can be drawn as follows:



Here,  $N = 80 \Rightarrow \frac{N}{2} = 40$ .

Mark the point A whose ordinate is 40 and its x-coordinate is 76.



Thus, median of the data is 76.


32.

**Sol:**

The frequency distribution table of more than type is as follows:

Marks (upper class limits)	Cumulative frequency (cf)
More than 0	$96 + 4 = 100$
More than 10	$90 + 6 = 96$
More than 20	$80 + 10 = 90$
More than 30	$70 + 10 = 80$
More than 40	$45 + 25 = 70$
More than 50	$23 + 22 = 45$
More than 60	$18 + 5 = 23$
More than 70	5

Taking lower class limits of on x-axis and their respective cumulative frequencies on y-axis, its ogive can be drawn as follows:




33.

**Sol:**

The frequency distribution table of more than type is as follows:

Height (in cm) (lower class limit)	Cumulative frequency (cf)
More than 135	$5 + 45 = 50$
More than 140	$8 + 37 = 45$
More than 145	$9 + 28 = 37$
More than 150	$12 + 16 = 28$
More than 155	$14 + 2 = 16$
More than 160	2

Taking lower class limits of on x-axis and their respective cumulative frequencies on y-axis, its ogive can be drawn as follows:




34.

**Sol:**

The frequency distribution table of more than type is as follows:

Height (in cm) (lower class limit)	Cumulative frequency (cf)
More than 140	$3 + 153 = 156$
More than 160	$8 + 145 = 153$
More than 180	$15 + 130 = 145$
More than 200	$40 + 90 = 130$
More than 220	$50 + 40 = 90$
More than 240	$30 + 10 = 40$
More than 260	10

Taking the lower class limits of on x-axis and their respective cumulative frequencies on y-axis, its ogive can be drawn as follows:






35.

**Sol:**

The frequency distribution table of more than type is as follows:


Production yield (kg/ha) ( lower class limits)	Cumulative frequency (cf)
More than 50	$2 + 98 = 100$
More than 55	$8 + 90 = 98$
More than 60	$12 + 78 = 90$
More than 65	$24 + 54 = 78$
More than 70	$38 + 16 = 54$
More than 75	16

Taking the lower class limits on x-axis and their respective cumulative on y-axis, its ogive can be drawn as follows:



Here,  $N = 100 \Rightarrow \frac{N}{2} = 50$ .

Mark the point A whose ordinate is 50 and its x-coordinate is 70.5.



Thus, median of the data is 70.5.


36.

**Sol:**

The frequency distribution table of less than type is as follows:

Weekly expenditure (in ₹) ( upper class limits)	Cumulative frequency (cf)
Less than 200	5
Less than 300	$5 + 6 = 11$
Less than 400	$11 + 11 = 22$
Less than 500	$22 + 13 = 35$
Less than 600	$35 + 5 = 40$
Less than 700	$40 + 4 = 44$
Less than 800	$44 + 3 = 47$
Less than 900	$47 + 2 = 49$

Taking the lower class limits on x-axis and their respective cumulative frequencies on y-axis, its ogive can be obtained as follows




Now,

The frequency distribution table of more than type is as follows:

Weekly expenditure (in ₹) ( lower class limits)	Cumulative frequency (cf)
More than 100	$44 + 5 = 49$
More than 200	$38 + 6 = 44$
More than 300	$27 + 11 = 38$
More than 400	$14 + 13 = 27$
More than 500	$9 + 5 = 14$
More than 600	$5 + 4 = 9$
More than 700	$2 + 3 = 5$
More than 800	2

Taking the lower class limits on x-axis and their respective cumulative frequencies on y-axis, its ogive can be obtained as follows:



37.


Sol:

From the given table, we may prepare than 'more than' frequency table as shown below:

Score	Number of candidates
More than 750	34
More than 700	52
More than 650	79
More than 600	103
More than 550	135
More than 500	175
More than 450	210
More than 400	230

We plot the points A(750, 34), B(700,52), C(650, 79), D(600, 103), E(550, 135), F(500, 175), G(450, 210) and H(400, 230).

Join AB, BC, CD, DE, EF, FG, GH and HA with a free hand to get the curve representing the 'more than type' series.



Here,  $N = 230$

$$\Rightarrow \frac{N}{2} = 115$$

From P (0, 115), draw PQ meeting the curve at Q. Draw QM meeting at M.

Clearly,  $OM = 590$  units

Hence, median = 590 units.

38.

**Sol:**

(i) From the given table, we may prepare the 'less than' frequency table as shown below:

Marks	Number of students
Less than 5	2
Less than 10	7
Less than 15	13
Less than 20	21
Less than 25	31
Less than 30	56
Less than 35	76
Less than 40	94
Less than 45	98
Less than 50	100

We plot the points A(5, 2), B(10, 7), C(15, 13), D(20, 21), E(25, 31), F(30, 56), G(35, 76) and H(40, 94), I(45, 98) and J(50, 100).


Join AB, BC, CD, DE, EF, FG, GH, HI, IJ and JA with a free hand to get the curve representing the 'less than type' series.

(ii) More than series:

Marks	Number of students
More than 0	100
More than 5	98
More than 10	93
More than 15	87
More than 20	79
More than 25	69
More than 30	44
More than 35	24
More than 40	6
More than 45	2

Now, on the same graph paper, we plot the points (0, 100), (5, 98), (10, 94), (15, 76), (20, 56), (25, 31), (30, 21), (35, 13), (40, 6) and (45, 2).

Join with a free hand to get the 'more than type' series.



The two curves intersect at point L. Draw  $LM \perp OX$  cutting the x-axis at M.

Clearly,  $M = 29.5$

Hence, Median = 29.5

39.

**Sol:**

(i) Less than series:

Marks	Number of students
Less than 144	3
Less than 148	12
Less than 152	36
Less than 156	67
Less than 160	109
Less than 164	173
Less than 168	248

Less than 172	230
Less than 176	416
Less than 180	450

We plot the points A(144, 3), B(148, 12), C(152, 36), D(156, 67), E(160, 109), F(164, 173), G(168, 248) and H(172, 330), I(176, 416) and J(180, 450).


Join AB, BC, CD, DE, EF, FG, GH, HI, IJ and JA with a free hand to get the curve representing the 'less than type' series.

(ii) More than series:

Marks	Number of students
More than 140	450
More than 144	447
More than 148	438
More than 152	414
More than 156	383
More than 160	341
More than 164	277
More than 168	202
More than 172	120
More than 176	34

Now, on the same graph paper, we plot the points A<sub>1</sub>(140, 450), B<sub>1</sub>(144, 447), C<sub>1</sub>(148, 438), D<sub>1</sub>(152, 414), E<sub>1</sub>(156, 383), F<sub>1</sub>(160, 277), H<sub>1</sub>(168, 202), I<sub>1</sub>(172, 120) and J<sub>1</sub>(176, 34).

Join A<sub>1</sub>B<sub>1</sub>, B<sub>1</sub>C<sub>1</sub>, C<sub>1</sub>D<sub>1</sub>, D<sub>1</sub>E<sub>1</sub>, E<sub>1</sub>F<sub>1</sub>, F<sub>1</sub>G<sub>1</sub>, G<sub>1</sub>H<sub>1</sub>, H<sub>1</sub>I<sub>1</sub> and I<sub>1</sub>J<sub>1</sub> with a free hand to get the 'more than type' series.



The two curves intersect at point L. Draw LM ⊥ OX cutting the x-axis at M. Clearly, M = 166cm

Hence, median = 166cm

## Exercise 9F

1.

**Sol:**

To find median let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative frequency (cf)
0-10	4	4
10-20	4	8
20-30	8	16
30-40	10	26
40-50	12	38
50-60	8	46
60-70	4	50
Total	$N = \sum f_i = 50$	

Now,  $N = 50 \Rightarrow \frac{N}{2} = 25$ 

The cumulative frequency just greater than 25 is 26, and the corresponding class is 30-40.

Thus, the median class is 30-40.

2.

**Sol:**

Here the maximum class frequency is 27, and the class corresponding to this frequency is 40-50. So the modal class is 40-50.

Now,

Modal class = 40-50, lower limit (/) of modal class = 40.

Thus, lower limit (/) of modal class is 40.

3.

**Sol:**

Here the maximum class frequency is 30, and the class corresponding to the frequency is 150-200. So, the modal class is 150-200.



Also, class mark of the modal class is  $\frac{150+200}{2} = 175$ .

4.

**Sol:**

If the number of observations is odd, then the median is  $\left(\frac{n+1}{2}\right)$ th observation.

Thus,  $\left(\frac{25+1}{2}\right) = 13$ th observation represents the median.

5.

**Sol:**

There is an empirical relationship between the three measures of central tendency:

$$3\text{median} = \text{mode} + 2\text{Mean}$$

$$\Rightarrow \text{Mean} = \frac{3\text{Median} - \text{Mode}}{2}$$

$$= \frac{3(1250) - 1000}{2}$$

$$= 1375$$

Thus, the mean is 1375.

6.

**Sol:**

Here the maximum class frequency is 25, and the class corresponding to this frequency is 40-60.

So, the modal class is 40-60.

Now, to find the median class let us put the data in the table given below:

Marks Obtained	Number of students ( $f_i$ )	Cumulative frequency (cf)
0-20	4	4
20-40	6	10
40-60	25	35
60-80	10	45
80-100	5	50
Total	$N = \sum f_i = 50$	

Now,  $N = 50 \Rightarrow \frac{N}{2} = 25$ .

The cumulative frequency just greater than 25 is 35, and the corresponding class is 40-60.

Thus, the median class is 40-60.

7.

**Sol:**

$$\text{Class mark} = \frac{\text{Upper limit} + \text{Lower limit}}{2}$$

$$\therefore \text{class mark of } 10-25 = \frac{10+25}{2}$$

$$= 17.5$$

$$\text{And class mark of } 35-55 = \frac{35+55}{2}$$

$$= 45$$

8.

**Sol:**

According to assumed-mean method,

$$\bar{x} = A + \frac{\sum f_i d_i}{\sum f_i}$$

$$= 25 + \frac{110}{50}$$

$$= 25 + 2.2$$

$$= 27.2$$

Thus, mean is 27.2.

9.

**Sol:**

According to the question,

$$4 = \frac{X}{36} \text{ and } 3 = \frac{Y}{64}$$

$$\Rightarrow X = 4 \times 36 \text{ and } Y = 3 \times 64$$

$$\Rightarrow X = 144 \text{ and } Y = 192$$

$$\text{Now, } X + Y = 144 + 192 = 336$$

$$\text{And total number of observations} = 36 + 64 = 100$$

$$\text{Thus, mean} = \frac{336}{100} = 3.36.$$

10.

**Sol:**

$$\text{Upper class boundary} = \text{Lowest class boundary} + \text{width} \times \text{number of classes}$$

$$= 8.1 + 2.5 \times 12$$

$$= 8.1 + 30$$

$$= 38.1$$

Thus, upper class boundary of the highest class is 38.1.

11.

**Sol:**

If number of observations is even, then the median will be the average of  $\left(\frac{n}{2}\right)$ th and the  $\left(\frac{n}{2} + 1\right)$ th observations.

In the given case,  $n = 10 \Rightarrow \left(\frac{n}{2}\right) \text{th} = 5 \text{th}$  and  $\left(\frac{n}{2} + 1\right) \text{th} = 6 \text{th}$  observation.

$$\text{Thus, } 63 = \frac{x+(x+2)}{2}$$

$$\Rightarrow 126 = 2x + 2$$

$$\Rightarrow 124 = 2x$$

$$\Rightarrow x = 62$$

Thus, the value of x is 62.

12.

**Sol:**

Since, 8 is less than 30 and 32 is more than 30, so the middle value remains unchanged

Thus, the median of 21 observations taken together is 30.

13.

**Sol:**

Arranging the observations in ascending order, we have

$$\frac{x}{5}, \frac{x}{4}, \frac{x}{3}, \frac{x}{2}, x$$

Thus, the median is  $\frac{x}{3}$

$$\Rightarrow \frac{x}{3} = 8$$

$$\Rightarrow x = 3 \times 8$$

$$\Rightarrow x = 24$$

Thus, the value of x is 24.

14.

**Sol:**

Here the maximum class frequency is 23, and the class corresponding to this frequency is 12-15.

So, the modal class is 12.15.

Now, to find the cumulative frequency let us put the data in the table given below:

Class	Frequency ( $f_i$ )	Cumulative frequency ( $cf$ )
3-6	7	7
6-9	13	20
9-12	10	30
12-15	23	53
15-18	4	57
18-21	21	78
21-24	16	94
Total	$N = \sum f_i = 94$	

Thus, the cumulative frequency of the modal class is 53.

15.

**Sol:**

Here the maximum class frequency is 18, and the class corresponding to this frequency is 40-60.

So, the modal class is 40-60.

Now,

Modal class = 40-60, lower limit ( $l$ ) of modal class = 40, class size ( $h$ )=20,

Frequency ( $f_1$ ) of the modal class =18,

Frequency ( $f_0$ ) of class preceding the modal class =6,

Frequency ( $f_2$ ) of class succeeding the modal class = 10.

Now, let us substitute these values in the formula:

$$Mode = l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

$$= 40 + \left( \frac{18-6}{36-6-10} \right) \times 20$$

$$= 40 + \left( \frac{12}{20} \right) \times 20$$

$$= 40 + 12$$

$$= 52$$

Hence, the mode is 52.

16.

**Sol:**

A 'less than type' cumulative frequency distribution table is given below:

Age (in years)	Cumulative frequency ( $cf$ )
Less than 20	60
Less than 30	102
Less than 40	157
Less than 50	227
Less than 60	280
Less than 70	300

17.

**Sol:**

$$\text{Here, } p = 11 + 12 = 23$$

$$\text{And } 33 + q = 46$$

$$\Rightarrow q = 46 - 33$$

$$= 13$$

Thus,  $p$  is 23 and  $q$  is 13.

Now,

Here the maximum class frequency is 20, and the class corresponding to this frequency is 500-600.

So, the modal class is 500-600.

$$\text{Also, } \Sigma f = N = 80$$

$$\Rightarrow \frac{N}{2} = 40.$$

The cumulative frequency just greater than 40 is 46, and the corresponding class is 400-500.

Thus, the median class is 400-500.

18.

**Sol:**

The cumulative frequency distribution table of more than type is as follows:

Monthly consumption (in units) (lower class limits)	Cumulative frequency ( $cf$ )
More than 65	$60 + 4 = 64$
More than 85	$55 + 5 = 60$
More than 105	$42 + 13 = 55$
More than 125	$22 + 20 = 42$
More than 145	$8 + 14 = 22$
More than 165	8

19.

**Sol:**

The frequency distribution is as follows:

Life-time (in days)	Frequency (f)
0-50	7
50-100	14
100-150	31
150-200	27
200-250	12
250-300	9

20.

**Sol:**

(a) The frequency distribution into the continuous form is as follows:

Marks obtained (in per cent)	Number of students (f)
10.5-20.5	141
20.5-30.5	221
30.5-40.5	439
40.5-50.5	529

50.5-60.5	495
60.5-70.5	322
70.5-80.5	153

(b) Now, to find the median class let us put the data in the table given below:

Marks obtained (in percent)	Number of students (f)	Cumulative frequency (cf)
10.5-20.5	141	141
20.5-30.5	221	362
30.5-40.5	439	801
40.5-50.5	529	1330
50.5-60.5	495	1825
60.5-70.5	322	2147
70.5-80.5	153	2300

Now,  $N = 2300$

$$\Rightarrow \frac{N}{2} = 1150$$

The cumulative frequency just greater than 1150 is 1330, and the corresponding class is 40.5-50.5.

Thus, the median class is 40.5-50.5

Now, class mark =  $\frac{\text{upper class limit} + \text{lower class limit}}{2}$

$$\frac{40.5 + 50.5}{2} = \frac{91}{2} = 45.5$$

Thus, class mark of the median class is 45.5

(c) Here the maximum class frequency is 529, and the class corresponding to this frequency is 40.5-50.5.

So, the modal class is 40.5-50.5 and its cumulative frequency is 1330.

21.

**Sol:** The given data is shown as follows:

Class	Frequency (f)	Class mark ( $x_i$ )	$f_i x_i$
0 – 10	8	5	40
10 – 20	P	15	15p
20 – 30	12	25	300
30 – 40	13	35	455
40 – 50	10	45	450
Total	$\sum f_i = 43 + p$		$\sum f_i x_i = 1245 + 15p$

The mean of given data is given by  $x = \frac{\sum f_i x_i}{\sum f_i}$

$$\Rightarrow 27 = \frac{1245 + 15p}{43 + p}$$

$$\Rightarrow 1161 + 27p = 1245 + 15p$$

$$\Rightarrow 27p - 15p = 1245 - 1161$$

$$\Rightarrow 12p = 84$$

$$\Rightarrow p = 7$$

Thus, the value of p is 7.

22.

**Sol:**

Let the missing frequency be x.

To find the median let us put data in the table given below:

Age (in years)	Number of persons (f)	Cumulative frequency (cf)
0-10	5	5
10-20	25	30
20-30	X	30 + x
30-40	18	48 + x
40-50	7	55 + x

The given median is 24,

$\therefore$  the median class is 20-30.

$\therefore l = 20, h = 10, N = 55 + x, f = x$  and  $cf = 30$

$$\text{Median} = l + \left( \frac{\frac{N}{2} - cf}{f} \right) \times h$$

$$\Rightarrow 24 = 20 + \left( \frac{\frac{55+x}{2} - 30}{x} \right) \times 10$$

$$\Rightarrow 24 - 20 = \left( \frac{55+x-60}{2x} \right) \times 10$$

$$\Rightarrow 4 = \left( \frac{x-5}{2x} \right) \times 10$$

$$\Rightarrow 8x = 10x - 50$$

$$\Rightarrow 2x = 50$$

$$\Rightarrow x = 25$$

Thus, the missing frequency is 25.



---

**Multiple choice questions**

1.

**Answer:** (d) Standard Deviation**Sol:**

The standard deviation is a measure of dispersion. It is the action or process of distributing thing over a wide area (nothing about central location).

2.

**Answer:** (a) Mean**Sol:**

The mean cannot be determined graphically because the values cannot be summed.

3.

**Answer:** (a) Mean**Sol:**

Mean is influenced by extreme values.

4.

**Answer:** (c) a histogram**Sol:**

The mode of a frequency distribution can be obtained graphically from a histogram.

5.

**Answer:** (d) ogives**Sol:**

This because median of a frequency distribution is found graphically with the help of ogives.

6.

**Answer:** (b) Median**Sol:**

The cumulative frequency table is useful in determining the median.

7.

**Answer:** (b) Median**Sol:**

The abscissa of the point of intersection of the ‘less than type’ and that of the ‘more than type’ cumulative frequency curves of a grouped data gives its median.

8.

**Answer:** (b) 0**Sol:**We know that  $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$ 

$$\Rightarrow \bar{x} \sum f_i = \sum f_i x_i \quad \dots (i)$$

$$\text{Now, } \sum f_i (x_i - \bar{x}) = \sum f_i x_i - \bar{x} \sum f_i$$

$$\Rightarrow \sum f_i (x_i - \bar{x}) = \sum f_i x_i - \sum f_i x_i \quad [\text{Using (i)}]$$

$$\Rightarrow \sum f_i (x_i - \bar{x}) = 0$$

9.

$$\text{Answer: (b) } u_i = \frac{(x_i - A)}{h}$$

**Sol:**

$$u_i = \frac{(x_i - A)}{h}$$

10.

**Answer:** (c) midpoints of the classes

**Sol:**

The  $d'_i$ 's are the deviations from A of midpoints of the classes.

11.

**Answer:** (b) centred at the class marks of the classes

**Sol:**

While computing the mean of the group data, we assume that the frequencies are centred at the class marks of the classes.

12.

**Answer:** (b)  $\text{mode} = (3 \times \text{median}) - (2 \times \text{mean})$

**Sol:**

$\text{mode} = (3 \times \text{median}) - (2 \times \text{mean})$

13.

**Answer:** (c) 20.5

**Sol:**

The x- coordinate represents the median of the given data.

Thus, median of the given data is 20.5.

14.

**Answer:** (b) 315**Sol:**

The class having the maximum frequency is the modal class.

So, the modal class is 150 – 155 and its lower limit is 150.

Also,  $N = 60$ 

$$\Rightarrow \frac{N}{2} = 30$$

The cumulative frequency just more than 30 is 37 and its class is 160 – 165, whose upper limit is 165.

$$\therefore \text{Required sum} = (150 + 165) = 315$$

15.

**Answer:** (c) 30 – 40**Sol:**

The class 30 – 40 has the maximum frequency, i.e., 30.

So, the modal class is 30 – 40.

16.

$$\text{Answer: (b) } x_k + h \left\{ \frac{f_k - f_{k-1}}{2f_k - f_{k-1} - f_{k+1}} \right\}$$

**Sol:**

$$x_k + h \left\{ \frac{f_k - f_{k-1}}{2f_k - f_{k-1} - f_{k+1}} \right\}$$

17.

**Answer:** (a)  $l + \left\{ h \times \frac{\left(\frac{N}{2} - cf\right)}{f} \right\}$

**Sol:**

$$l \times \left\{ h \times \frac{\left(\frac{N}{2} - cf\right)}{f} \right\}$$

18.

**Answer:** (c) 9.2**Sol:**

It is given that the mean and median are 8.9 and 9, respectively,

$$\therefore \text{Mode} = (3 \times \text{Median}) - (2 \times \text{Mean})$$

$$\Rightarrow \text{Mode} = (3 \times 9) - (2 \times 8.9)$$

$$= 27 - 17.8$$

$$= 9.2$$

19.

**Answer:** (b) 57.5**Sol:**

Class interval	35 – 45	45 – 55	55 – 65	65 – 75
Frequency	8	12	20	10
Cumulative frequency	8	20	40	50

Here,  $N = 50$ 

$$\Rightarrow \frac{N}{2} = 25, \text{ which lies in the class interval of } 55 - 65.$$
Now,  $cf = 55$ ,  $f = 20$  and  $l = 50$ 

$$\therefore \text{Median} = l + \left\{ h \times \frac{\left(\frac{N}{2} - cf\right)}{f} \right\}$$

$$= 50 + \frac{65-55}{20} \times (25 - 20)$$

$$= 57.5$$

20.

**Answer:** (c) 24.4**Sol:**

The maximum frequency is 25 and the modal class is 22 - 26.

Now,  $x_k = 22$ ,  $f_k = 25$ ,  $f_{k-1} = 16$ ,  $f_{k+1} = 19$  and  $h = 4$ 

$$\begin{aligned} \therefore \text{Mode} &= x_k + h \left\{ \frac{f_k - f_{k-1}}{2f_k - f_{k-1} - f_{k+1}} \right\} \\ &= 22 + 4 \times \frac{(25-16)}{(2 \times 25 - 16 - 19)} \\ &= 22 + 4 \times \frac{(25-16)}{(50-16-19)} \\ &= 22 + 4 \times \frac{9}{15} \\ &= 22 + \frac{12}{5} \\ &= 22 + 2.4 \\ &= 24.4 \end{aligned}$$

21.

**Answer:** (c) 24**Sol:**

$$\text{Mode} = (3 \times \text{median}) - (2 \times \text{mean})$$

$$\Rightarrow (3 \times \text{median}) = (\text{mode} + 2 \text{ mean})$$

$$\Rightarrow (3 \times \text{median}) = 16 + 56$$

$$\Rightarrow (3 \times \text{median}) = 72$$

$$\Rightarrow \text{Median} = \frac{72}{3}$$

$$\therefore \text{Median} = 24$$

22.

**Answer:** (b) 24.5**Sol:**

$$\text{Mode} = (3 \times \text{median}) - (2 \times \text{mean})$$

$$\Rightarrow (2 \times \text{mean}) = (3 \times \text{median}) - \text{mode}$$

$$\Rightarrow (2 \times \text{mean}) = 3 \times 26 - 29$$

$$\Rightarrow (2 \times \text{mean}) = 49$$

$$\Rightarrow \text{Mean} = \frac{49}{2}$$

$$\therefore \text{Mean} = 24.5$$

23.

**Answer:** (c) mean = mode = median**Sol:**

A symmetric distribution is one where the left and right hand sides of the distribution are roughly equally balanced around the mean.

24.

**Answer:** (c) 13**Sol:**

Converting the given data into a frequency table, we get:

Monthly income	No. of families	Frequency
30,000 and above	15	15
25,000 – 30,000	37	$(37 - 15) = 22$
20,000 – 25,000	50	$(50 - 37) = 13$
18,000 – 20,000	69	$(69 - 50) = 19$
14,000 – 18,000	85	$(85 - 69) = 16$
10,000 – 14,000	100	$(100 - 85) = 15$

Hence, the number of families having an income range of Rs. 20,000 – Rs. 25,000 is 13.  
The correct option is (c).

25.

**Answer:** (b) 9**Sol:**

First 8 prime numbers are 2, 3, 5, 7, 11, 13, 17 and 19.

Median of 8 numbers is average of 4<sup>th</sup> and 5<sup>th</sup> terms.

i.e., average of 7 and 11  
Thus, the median is 9.

26.

**Answer:** (d) 19

**Sol:**

It is given that mean of 20 numbers is zero.

(a) e., average of 20 numbers is zero.

i.e., sum of 20 numbers is zero.

Thus, at most, there can be 19 positive numbers.

(such that if sum of 19 positive numbers is  $x$ , 20<sup>th</sup> number will be  $-x$ )

27.

**Answer:** (c) 15

**Sol:**

Median of 6 numbers is the average of 3<sup>rd</sup> and 4<sup>th</sup> term.

$$\therefore 13 = \frac{(x-1)+(x-3)}{2}$$

$$\Rightarrow 26 = 2x - 4$$

$$\Rightarrow 2x = 30$$

$$\Rightarrow x = 15$$

Thus,  $x$  is equal to 15.

28.

**Answer:** (c) -20

**Sol:**

$$\text{Mean} = \frac{\text{sum of observations}}{\text{number of observations}}$$

$$\Rightarrow 15 = \frac{2+7+6+x}{4}$$

$$\Rightarrow 60 = 15 + x$$

$$\Rightarrow x = 45 \quad \dots\dots(1)$$

Now,

$$\text{Mean} = \frac{\text{sum of observations}}{\text{number of observations}}$$

$$\Rightarrow 10 = \frac{18+1+6+x+y}{5}$$

$$\Rightarrow 50 = 25 + x + y$$

$$\Rightarrow y = 25 - x$$



$$\Rightarrow y = 25 - 45 \quad [\text{From (1)}]$$

$$\Rightarrow y = -20$$

29.

**Sol:**

Column I	Column II
(a) The most frequent value in a data is known as .....	(s) mode
(b) which of the following cannot be determined graphically out of mean, mode and median?	(r) mean
(c) An ogive is used to determine .....	(q) median
(d) out of mean, mode, median and standard deviation, which is not a measure of tendency?	(p) standard deviation

30.

**Sol:**

(a) Both Assertion (A) and reason (R) are true and Reason (R) is a correct explanation of Assertion (A).

Clearly, reason (R) is true.

Using the relation in reason (R), we have:

$$2 \text{ mean} = (3 \times \text{median}) - \text{mode} = (3 \times 150) - 154 = 450 - 154 = 296$$

$$\Rightarrow \text{Mean} = 148, \text{ which is true.}$$

$\therefore$  This assertion (A) and reason (R) are both true and reason (R) is the correct explanation of assertion (A).

31.

**Sol:**

(b) Both assertion (A) and reason (R) are true, but reason (R) is not a correct explanation of assertion (A).

Clearly, reason (R) is true.

The maximum frequency is 23 and the modal class is 12 – 15.

Now,  $x_k = 12$ ,  $f_k = 23$ ,  $f_{k-1} = 21$ ,  $f_{k+1} = 23$  and  $h = 3$

$$\therefore \text{Mode} = \left\{ 12 + 3 \times \frac{(23-21)}{(2 \times 23 - 21 - 10)} \right\}$$

$$= \left( 12 + 3 \times \frac{2}{15} \right)$$

$$= (12 + 0.4)$$

$$= 12$$

$\therefore$  Assertion (A) is true.

However, reason (R) is not a correct explanation of assertion (A).